

β -Arrestin 1 (D7Z3W) XP[®] Rabbit mAb

Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications:	Reactivity:	Sensitivity:	MW (kDa):	Source/Isotype:	UniProt ID:	Entrez-Gene Id:
W, IHC-P	H M R	Endogenous	51	Rabbit IgG	#P49407	408

Product Usage Information**Application**

Western Blotting
Immunohistochemistry (Paraffin)

Dilution

1:1000
1:300

Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 μ g/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at -20°C . Do not aliquot the antibody.

For a carrier free (BSA and azide free) version of this product see product #47698.

Specificity/Sensitivity

β -Arrestin (D7Z3W) XP[®] Rabbit mAb recognizes endogenous levels of total β -arrestin 1 protein. This antibody does not cross-react with other arrestin proteins.

Source / Purification

Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to residues surrounding Arg363 of human β -arrestin 1 protein.

Background

Arrestin proteins function as negative regulators of G protein-coupled receptor (GPCR) signaling. Cognate ligand binding stimulates GPCR phosphorylation, which is followed by binding of arrestin to the phosphorylated GPCR and the eventual internalization of the receptor and desensitization of GPCR signaling (1). Four distinct mammalian arrestin proteins are known. Arrestin 1 (also known as S-arrestin) and arrestin 4 (X-arrestin) are localized to retinal rods and cones, respectively. Arrestin 2 (also known as β -arrestin 1) and arrestin 3 (β -arrestin 2) are ubiquitously expressed and bind to most GPCRs (2). β -arrestins function as adaptor and scaffold proteins and play important roles in other processes, such as recruiting c-Src family proteins to GPCRs in Erk activation pathways (3,4). β -arrestins are also involved in some receptor tyrosine kinase signaling pathways (5-8). Additional evidence suggests that β -arrestins translocate to the nucleus and help regulate transcription by binding transcriptional cofactors (9,10).

The non-visual β -arrestins respond to glucocorticoid signaling, with differential responses observed among family members. Specifically, β -arrestin 1 expression is increased in response to glucocorticoid receptor activation whereas β -arrestin 2 shows a concomitant decrease in expression (11).

Background References

- Shenoy, S.K. and Lefkowitz, R.J. (2005) *Sci STKE* 2005, cm10.
- Lefkowitz, R.J. and Shenoy, S.K. (2005) *Science* 308, 512-7.
- Luttrell, L.M. et al. (1999) *Science* 283, 655-61.
- Luttrell, L.M. et al. (1999) *Curr Opin Cell Biol* 11, 177-83.
- Luttrell, L.M. and Lefkowitz, R.J. (2002) *J Cell Sci* 115, 455-65.
- Waters, C. et al. (2004) *Semin Cell Dev Biol* 15, 309-23.
- Lefkowitz, R.J. and Whalen, E.J. (2004) *Curr Opin Cell Biol* 16, 162-8.
- Waters, C.M. et al. (2005) *Cell Signal* 17, 263-77.
- Kang, J. et al. (2005) *Cell* 123, 833-47.
- Ma, L. and Pei, G. (2007) *J Cell Sci* 120, 213-8.
- Oakley, R.H. et al. (2012) *Proc Natl Acad Sci U S A* 109, 17591-6.

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v nonfat dry milk, 1X TBS, 0.1% Tween[®] 20 at 4°C with gentle shaking, overnight.

Applications Key

W: Western Blotting **IHC-P:** Immunohistochemistry (Paraffin)

Cross-Reactivity Key

H: Human **M:** Mouse **R:** Rat

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

SignalStain is a registered trademark of Cell Signaling Technology, Inc.

XP is a registered trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.