


<b>#3024</b> Store at -20C	<b>Phospho-IGF-I Receptor <math>\beta</math></b> <b>(Tyr1135/1136)/Insulin Receptor <math>\beta</math></b> <b>(Tyr1150/1151) (19H7) Rabbit mAb</b>	
	<b>Orders:</b> 877-616-CELL (2355) orders@cellsignal.com  <b>Support:</b> 877-678-TECH (8324)  <b>Web:</b> info@cellsignal.com cellsignal.com	<b>3 Trask Lane   Danvers   Massachusetts   01923   USA</b>

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications:	Reactivity:	Sensitivity:	MW (kDa):	Source/Isotype:	UniProt ID:	Entrez-Gene ID:
W, W-S	H M R	Endogenous	95	Rabbit IgG	#P06213, #P08069	3643, 3480

#### Product Usage Information

#### Application

Western Blotting  
Simple Western™

#### Dilution

1:1000  
1:10 - 1:50

#### Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100  $\mu$ g/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at -20°C. Do not aliquot the antibody.

#### Specificity/Sensitivity

Phospho-IGF-I Receptor  $\beta$  (Tyr1135/1136)/Insulin Receptor  $\beta$  (Tyr1150/1151) (19H7) Rabbit mAb detects endogenous levels of IGF-I receptor and insulin receptor only when phosphorylated at tyrosine 1135/1136 or tyrosine 1150/1151, respectively. It does not cross-react with other related tyrosine-phosphorylated tyrosine kinases.

#### Species predicted to react based on 100% sequence homology

Bovine, Dog

#### Source / Purification

Monoclonal antibody is produced by immunizing animals with a synthetic phosphopeptide corresponding to residues surrounding Tyr1135/1136 of human IGF-I receptor  $\beta$ .

#### Background

Type I insulin-like growth factor receptor (IGF-IR) is a transmembrane receptor tyrosine kinase that is widely expressed in many cell lines and cell types within fetal and postnatal tissues (1-3). Receptor autophosphorylation follows binding of the IGF-I and IGF-II ligands. Three tyrosine residues within the kinase domain (Tyr1131, Tyr1135, and Tyr1136) are the earliest major autophosphorylation sites (4). Phosphorylation of these three tyrosine residues is necessary for kinase activation (5,6). Insulin receptors (IRs) share significant structural and functional similarity with IGF-I receptors, including the presence of an equivalent tyrosine cluster (Tyr1146/1150/1151) within the kinase domain activation loop. Tyrosine autophosphorylation of IRs is one of the earliest cellular responses to insulin stimulation (7). Autophosphorylation begins with phosphorylation at Tyr1146 and either Tyr1150 or Tyr1151, while full kinase activation requires triple tyrosine phosphorylation (8).

#### Background References

- Adams, T.E. et al. (2000) *Cell Mol Life Sci* 57, 1050-93.
- Baserga, R. (2000) *Oncogene* 19, 5574-81.
- Scheidegger, K.J. et al. (2000) *J Biol Chem* 275, 38921-8.
- Hernández-Sánchez, C. et al. (1995) *J Biol Chem* 270, 29176-81.
- Lopaczynski, W. et al. (2000) *Biochem Biophys Res Commun* 279, 955-60.
- Baserga, R. (1999) *Exp Cell Res* 253, 1-6.
- White, M.F. et al. (1985) *J Biol Chem* 260, 9470-8.
- White, M.F. et al. (1988) *J Biol Chem* 263, 2969-80.

#### Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

#### Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v nonfat dry milk, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

#### Applications Key

**W:** Western Blotting **W-S:** Simple Western™

#### Cross-Reactivity Key

**H:** Human **M:** Mouse **R:** Rat

#### Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit [cellsignal.com/trademarks](http://cellsignal.com/trademarks) for more information.

#### Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's

terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.

**Orders: 877-616-CELL (2355) • orders@cellsignal.com • Support: 877-678-TECH (8324) • info@cellsignal.com • Web: cellsignal.com**  
For Research Use Only. Not for Use in Diagnostic Procedures.