



Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

Store at -20C
#3043

MAPKAPK-3 Antibody

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications:	Reactivity:	Sensitivity:	MW (kDa):	Source/Isotype:	UniProt ID:	Entrez-Gene Id:
W	H R Mk	Endogenous	42	Rabbit	#Q16644	7867

Product Usage Information

Application

Western Blotting

Dilution

1:1000

Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA and 50% glycerol. Store at -20°C. Do not aliquot the antibody.

Specificity/Sensitivity

MAPKAPK-3 Antibody detects endogenous levels of total MAPKAPK-3 protein. This antibody does not cross-react with MAPKAPK-2 or -5.

Source / Purification

Polyclonal antibodies are produced by immunizing animals with a synthetic peptide corresponding to the amino-terminal region of human MAPKAPK-3. Antibodies are purified by protein A and peptide affinity chromatography.

Background

MAPKAPK-3 has a single potential SH3-binding site in the proline-rich amino terminus, a putative ATP-binding site, 2 MAP kinase phosphorylation site motifs, and a putative nuclear localization signal. It shares 72% nucleotide and 75% amino acid identity with MAPKAPK-2 (1). MAPKAPK-3 has been shown to be activated by growth inducers and stress stimulation of cells. *In vitro* studies have demonstrated that Erk, p38 MAP kinase, and Jun amino-terminal kinase are able to phosphorylate and activate MAPKAPK-3, which suggested a role for this kinase as an integrative element of signaling in both mitogen and stress responses (2). MAPKAPK-3 was reported to interact with, phosphorylate, and repress the activity of E47, which is a basic helix-loop-helix transcription factor involved in the regulation of tissue-specific gene expression and cell differentiation (3). MAPKAPK-3 may also support luteal maturation through the phosphorylation and activation of the nuclear transcription factor CREB (4).

Background References

- Sithanandam, G. et al. (1996) *Mol. Cell Biol.* 16, 868-876.
- Ludwig, S. et al. (1996) *Mol. Cell Biol.* 16, 6687-6697.
- Neufeld, B. et al. (2000) *J. Biol. Chem.* 275, 20239-20242.
- Maizels, E.T. et al. (2001) *Mol. Endocrinol.* 15, 716-733.

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

Applications Key

W: Western Blotting

Cross-Reactivity Key

H: Human **R:** Rat **Mk:** Monkey

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in

any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.