


#3104 Store at -20C	Phospho-SMAD2 (Ser245/250/255) Antibody	
	Orders: 877-616-CELL (2355) orders@cellsignal.com	
	Support: 877-678-TECH (8324)	
	Web: info@cellsignal.com cellsignal.com	
3 Trask Lane Danvers Massachusetts 01923 USA		

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications:	Reactivity:	Sensitivity:	MW (kDa):	Source/Isotype:	UniProt ID:	Entrez-Gene ID:
W	H M R Mk	Endogenous	60	Rabbit	#Q15796	4087

Product Usage Information

Application

Western Blotting

Dilution

1:1000

Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA and 50% glycerol. Store at -20°C. Do not aliquot the antibody.

Specificity/Sensitivity

Phospho-SMAD2 (Ser245/250/255) Antibody detects endogenous levels of SMAD2 only when phosphorylated at serines 245, 250 or 255.

Source / Purification

Polyclonal antibodies are produced by immunizing animals with a synthetic phosphopeptide corresponding to residues surrounding serines 245/250/255 of human SMAD2. Antibodies are purified by protein A and peptide affinity chromatography.

Background

Members of the SMAD family of signal transduction molecules are components of a critical intracellular pathway that transmit TGF-β signals from the cell surface into the nucleus. Three distinct classes of SMADs have been defined: the receptor-regulated SMADs (R-SMADs), which include SMAD1, 2, 3, 5, and 9; the common-mediator SMAD (co-SMAD), SMAD4; and the antagonistic or inhibitory SMADs (I-SMADs), SMAD6 and 7 (1-5). Activated type I receptors associate with specific R-SMADs and phosphorylate them on a conserved carboxy-terminal SSXS motif. The phosphorylated R-SMADs dissociate from the receptor and form a heteromeric complex with SMAD4, initiating translocation of the heteromeric SMAD complex to the nucleus. Once in the nucleus, SMADs recruit a variety of DNA binding proteins that function to regulate transcriptional activity (6-8).

Oncogenic Ras antagonizes TGF-beta signaling and inhibits the nuclear accumulation of Smad2 and Smad3, which may be explained through MAP kinase dependent phosphorylation of these Smads (9). Cell stimulation with EGF leads to phosphorylation of Smad2 at a cluster of serine-proline sites within its linker region, including Ser245, 250, and 255 (9).

Background References

- Heldin, C.H. et al. (1997) *Nature* 390, 465-71.
- Attisano, L. and Wrana, J.L. (1998) *Curr Opin Cell Biol* 10, 188-94.
- Derynck, R. et al. (1998) *Cell* 95, 737-40.
- Massagué, J. (1998) *Annu Rev Biochem* 67, 753-91.
- Whitman, M. (1998) *Genes Dev* 12, 2445-62.
- Wrana, J.L. (2000) *Sci STKE* 2000, re1.
- Attisano, L. and Wrana, J.L. (2002) *Science* 296, 1646-7.
- Moustakas, A. et al. (2001) *J Cell Sci* 114, 4359-69.
- Kretzschmar, M. et al. (1999) *Genes Dev* 13, 804-16.

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

Applications Key

W: Western Blotting

Cross-Reactivity Key

H: Human **M:** Mouse **R:** Rat **Mk:** Monkey

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless

separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.

Orders: 877-616-CELL (2355) • orders@cellsignal.com • Support: 877-678-TECH (8324) • info@cellsignal.com • Web: cellsignal.com
For Research Use Only. Not for Use in Diagnostic Procedures.