



**Orders:** 877-616-CELL (2355)  
orders@cellsignal.com

**Support:** 877-678-TECH (8324)

**Web:** info@cellsignal.com  
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

Store at -20C  
#3130

## c-IAP2 (58C7) Rabbit mAb

For Research Use Only. Not for Use in Diagnostic Procedures.

<b>Applications:</b> W, W-S, IP	<b>Reactivity:</b> H	<b>Sensitivity:</b> Endogenous	<b>MW (kDa):</b> 70	<b>Source/Isotype:</b> Rabbit IgG	<b>UniProt ID:</b> #Q13489	<b>Entrez-Gene Id:</b> 330
------------------------------------	-------------------------	-----------------------------------	------------------------	--------------------------------------	-------------------------------	-------------------------------

### Product Usage Information

#### Application

Western Blotting  
Simple Western™  
Immunoprecipitation

#### Dilution

1:1000  
1:50 - 1:250  
1:100

### Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at -20°C. Do not aliquot the antibody.

### Specificity/Sensitivity

c-IAP2 (58C7) Rabbit mAb detects endogenous levels of total c-IAP2 protein. No cross reactivity was detected with other family members.

### Species predicted to react based on 100% sequence homology

Monkey

### Source / Purification

Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to residues surrounding Val117 of human c-IAP2.

### Background

The inhibitor of apoptosis protein (IAP) family consists of an evolutionarily conserved group of apoptosis inhibitors containing a conserved 70 amino acid BIR (baculovirus inhibitor repeat) domain (1,2). Human members of this family include c-IAP1, c-IAP2, XIAP, survivin, livin, and NAIP. Overexpression of IAP family members, particularly survivin and livin, in cancer cell lines and primary tumors suggests an important role for these proteins in cancer progression (3-5). In general, the IAP proteins function through direct interactions to inhibit the activity of several caspases, including caspase-3, caspase-7, and caspase-9 (5,6). In addition, binding of IAP family members to the mitochondrial protein Smac blocks their interaction with caspase-9, thereby allowing the processing and activation of the caspase (2).

### Background References

1. Deveraux, Q.L. and Reed, J.C. (1999) *Genes Dev* 13, 239-52.
2. Deveraux, Q.L. et al. (1998) *EMBO J* 17, 2215-23.
3. Altieri, D.C. et al. (1999) *Lab Invest* 79, 1327-33.
4. Tamm, I. et al. (2000) *Clin Cancer Res* 6, 1796-803.
5. Kasof, G.M. and Gomes, B.C. (2001) *J Biol Chem* 276, 3238-46.
6. Deveraux, Q.L. et al. (1997) *Nature* 388, 300-4.

### Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

### Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween@ 20 at 4°C with gentle shaking, overnight.

### Applications Key

**W:** Western Blotting **W-S:** Simple Western™ **IP:** Immunoprecipitation

### Cross-Reactivity Key

**H:** Human

### Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit [cellsignal.com/trademarks](http://cellsignal.com/trademarks) for more information.

### Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless

separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.