



Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

Store at -20C
#3159

IFN- γ (3F1E3) Mouse mAb

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications: W, IP, E-P	Reactivity: H	Sensitivity: Recombinant protein	MW (kDa): 17	Source/Isotype: Mouse IgG1	UniProt ID: #P01579	Entrez-Gene Id: 3458
------------------------------------	-------------------------	--	------------------------	--------------------------------------	-------------------------------	--------------------------------

Product Usage Information

Application

Western Blotting
Immunoprecipitation
Peptide ELISA (DELFI A)

Dilution

1:1000
1:50
1:100

Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 μ g/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at -20°C . Do not aliquot the antibody.

Specificity/Sensitivity

IFN- γ (3F1E3) Mouse mAb detects recombinant human IFN- γ .

Source / Purification

Monoclonal antibody is produced by immunizing animals with Ni-NTA purified recombinant human IFN- γ expressed in *E. Coli*. Antibodies were prepared from ascites.

Background

IFN- γ plays key roles in both the innate and adaptive immune response. IFN- γ activates the cytotoxic activity of innate immune cells, such as macrophages and NK cells (1,2). IFN- γ production by NK cells and antigen presenting cells (APCs) promotes cell-mediated adaptive immunity by inducing IFN- γ production by T lymphocytes, increasing class I and class II MHC expression, and enhancing peptide antigen presentation (1). Due to differences in the degree of glycosylation, there are three forms of IFN- γ , with approximate molecular weights of 25, 20, and 15.5 kDa by SDS-PAGE (5). The anti-viral activity of IFN- γ is due to its induction of PKR and other regulatory proteins. Binding of IFN- γ to the IFNGR1/IFNGR2 complex promotes dimerization of the receptor complexes to form the (IFNGR1/IFNGR2)₂-IFN- γ dimer. Binding induces a conformational change in receptor intracellular domains and signaling involves Jak1, Jak2, and Stat1 (3). The critical role of IFN- γ in amplification of immune surveillance and function is supported by increased susceptibility to pathogen infection by IFN- γ or IFNGR knockout mice and in humans with inactivating mutations in *IFNGR1* or *IFNGR2*. IFN- γ also appears to have a role in atherosclerosis (4).

IFN- γ , also known as type II interferon, is produced mainly in activated T lymphocytes and natural killer cells (4) and has broad effects on various cells of the immune system. Many signaling proteins including IL-2, FGF, and EGF induce the synthesis of IFN- γ .

Background References

- Schroder, K. et al. (2004) *J Leukoc Biol* 75, 163-89.
- Martinez, F.O. et al. (2009) *Annu Rev Immunol* 27, 451-83.
- Kotenko, S.V. et al. (1995) *J Biol Chem* 270, 20915-21.
- McLaren, J.E. and Ramji, D.P. (2009) *Cytokine Growth Factor Rev* 20, 125-35.
- Kelker, H.C. et al. (1984) *J Biol Chem* 259, 4301-4.
- Young, H.A. and Hardy, K.J. (1995) *J Leukoc Biol* 58, 373-81.

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v nonfat dry milk, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

Applications Key

W: Western Blotting **IP:** Immunoprecipitation **E-P:** Peptide ELISA (DELFI A)

Cross-Reactivity Key

H: Human

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.