



Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

CLK3 Antibody

#3256 Store at -20C

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications:	Reactivity:	Sensitivity:	MW (kDa):	Source/Isotype:	UniProt ID:	Entrez-Gene Id:
W, IF-IC	H M R Mk	Endogenous	59, 17	Rabbit	#P49761	1198

Product Usage Information

Application

Western Blotting
Immunofluorescence (Immunocytochemistry)

Dilution

1:1000
1:25

Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA and 50% glycerol. Store at -20°C. Do not aliquot the antibody.

Specificity/Sensitivity

CLK3 Antibody detects endogenous levels of full-length and truncated forms of CLK3 protein.

Source / Purification

Polyclonal antibodies are produced by immunizing animals with a synthetic peptide corresponding to residues surrounding Tyr61 of human CLK3 (P49761-1). Antibodies were purified by protein A and peptide affinity chromatography.

Background

The cdc2-like kinase (CLK) family contains at least four highly conserved isoforms: CLK1, CLK2, CLK3 and CLK4 (1,2). CLKs are dual specificity kinases that autophosphorylate on serine, threonine and tyrosine residues and phosphorylate exogenous substrates on serine and threonine residues (2). CLK family members exist as both a full-length catalytically active form and an alternatively-spliced, inactive truncated form (1). A family of highly phosphorylated proteins, called serine and arginine rich (SR) proteins, are phosphorylated by CLKs (3-5). SR proteins are splicing factors that regulate the assembly of the spliceosome, a macromolecular complex where RNA splicing occurs in the nucleus. They are also involved in the selection of splice sites. Thus, CLKs may play important roles in regulating RNA splicing. CLK3 is abundantly expressed in the testis and, similar to other family members, has been implicated in regulating RNA splicing (6-8).

Background References

- Hanes, J. et al. (1994) *J. Mol. Biol.* 244, 665-672.
- Nayler, O. et al. (1997) *Biochem. J.* 326, 693-700.
- Colwill, K. et al. (1996) *EMBO J.* 15, 265-275.
- Prasad, J. and Manley, J.L. (2003) *Mol. Cell Biol.* 23, 4139-4149.
- Muraki, M. et al. (2004) *J. Biol. Chem.* 279, 24246-24254.
- Menegay, H. et al. (1999) *Exp. Cell Res.* 253, 463-473.
- Becker, W. et al. (1996) *Biochim. Biophys. Acta* 1312, 63-67.
- Duncan, P.I. et al. (1998) *Exp. Cell Res.* 241, 300-308.

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

Applications Key

W: Western Blotting **IF-IC:** Immunofluorescence (Immunocytochemistry)

Cross-Reactivity Key

H: Human **M:** Mouse **R:** Rat **Mk:** Monkey

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

Alexa Fluor is a registered trademark of Life Technologies Corporation.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no

force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.