

## 8769

## Axl (L283) Antibody



Orders: 877-616-CELL (2355)

orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com

cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

## For Research Use Only. Not for Use in Diagnostic Procedures.

Applications: W	Reactivity: H Mk	<b>Sensitivity:</b> Endogenous	<b>MW (kDa):</b> 138	Source/Isotype: Rabbit	UniProt ID: #P30530	Entrez-Gene Id: 558
Product Usage Information		Application Dilution				
		Western Blotting			1:1000	
Storage		Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 μg/ml BSA and 50% glycerol. Store at – 20°C. Do not aliquot the antibody.				
Specificity/Sensitivity		Axl (L283) Antibody detects endogenous levels of total Axl protein.				
Source / Purification		Polyclonal antibodies are produced by immunizing animals with a synthetic peptide corresponding to sequences surrounding amino acid residue Leu283 of the human Axl extracellular domain. Antibodies are purified by protein A and peptide affinity chromatography.				
Background		Axl, Sky, and Mer are three members of a receptor tyrosine kinase (RTK) family that share a conserved intracellular tyrosine kinase domain and an extracellular domain similar to those seen in cell adhesion molecules. These RTKs bind the vitamin K-dependent protein growth-arrest-specific gene 6 (Gas6), which is structurally related to the protein S anticoagulation factor (1). Upon binding to its receptor, Gas6 activates phosphatidylinositol 3-kinase (PI3K) and its downstream targets Akt and S6K, as well as NF-kB (2,3). A large body of evidence supports a role for Gas6/Axl signaling in cell growth and survival in normal and cancer cells (4).				
Background References		1. Crosier, K.E. and Crosier, P.S. (1997) <i>Pathology</i> 29, 131-5. 2. Demarchi, F. et al. (2001) <i>J Biol Chem</i> 276, 31738-44. 3. Lee, W.P. et al. (2002) <i>Oncogene</i> 21, 329-36. 4. Bellosta, P. et al. (1997) <i>Oncogene</i> 15, 2387-97.				
Species Reactivity		Species reactivity is determined by testing in at least one approved application (e.g., western blot).				
Western Blot Buffer		IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.				
Applications Key		W: Western Blotting				
Cross-Reactivity Key		H: Human Mk: Monkey				
Trademarks and Patents		Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.				
		All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.				
Limited Uses		Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.				
		Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for				

Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party,

whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.