



**Orders:** 877-616-CELL (2355)  
orders@cellsignal.com

**Support:** 877-678-TECH (8324)

**Web:** info@cellsignal.com  
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

Store at -20C  
#3299

## Annexin A1 Antibody

For Research Use Only. Not for Use in Diagnostic Procedures.

<b>Applications:</b> W	<b>Reactivity:</b> H M R Mk	<b>Sensitivity:</b> Endogenous	<b>MW (kDa):</b> 37	<b>Source/Isotype:</b> Rabbit	<b>UniProt ID:</b> #P04083	<b>Entrez-Gene Id:</b> 301
---------------------------	--------------------------------	-----------------------------------	------------------------	----------------------------------	-------------------------------	-------------------------------

### Product Usage Information

#### Application

Western Blotting

#### Dilution

1:1000

### Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA and 50% glycerol. Store at -20°C. Do not aliquot the antibody.

### Specificity/Sensitivity

Annexin A1 Antibody detects endogenous levels of total annexin A1 protein.

### Source / Purification

Polyclonal antibodies are produced by immunizing animals with a synthetic peptide corresponding to human annexin A1. Antibodies are purified by peptide affinity chromatography.

### Background

The annexin superfamily consists of 13 calcium or calcium and phospholipid binding proteins with high biological and structural homology (1). Annexin-1 (ANXA1) is the first characterized member of the annexin family of proteins and is able to bind to cellular membranes in a calcium-dependent manner, promoting membrane fusion and endocytosis (2-4). Annexin A1 has anti-inflammatory properties and inhibits phospholipase A2 activity (5,6). Annexin A1 can accumulate on internalized vesicles after EGF-stimulated endocytosis and may be required for a late stage in inward vesiculation (7). Phosphorylation by PKC, EGFR, and Chak1 results in inhibition of annexin A1 function (8-10). Annexin A1 has also been identified as one of the 'eat-me' signals on apoptotic cells that are to be recognized and ingested by phagocytes (11). Annexin A1, as an endogenous anti-inflammatory mediator, has roles in many diverse cellular functions, such as membrane aggregation, inflammation, phagocytosis, proliferation, apoptosis, and tumorigenesis and cancer development (12-14).

### Background References

1. Raynal, P. and Pollard, H.B. (1994) *Biochim Biophys Acta* 1197, 63-93.
2. Blackwell, G.J. et al. (1980) *Nature* 287, 147-9.
3. Rothhut, B. et al. (1983) *Biochem Biophys Res Commun* 117, 878-84.
4. Hirata, F. et al. (1981) *Proc Natl Acad Sci USA* 78, 3190-4.
5. Kim, K.M. et al. (1994) *FEBS Lett* 343, 251-5.
6. Kim, S.W. et al. (2001) *J Biol Chem* 276, 15712-9.
7. White, I.J. et al. (2006) *EMBO J* 25, 1-12.
8. Varticovski, L. et al. (1988) *Biochemistry* 27, 3682-90.
9. Dorovkov, M.V. and Ryazanov, A.G. (2004) *J Biol Chem* 279, 50643-6.
10. Wang, W. and Creutz, C.E. (1994) *Biochemistry* 33, 275-82.
11. Arur, S. et al. (2003) *Dev Cell* 4, 587-98.
12. Perretti, M. and Gavins, F.N. (2003) *News Physiol Sci* 18, 60-4.
13. Parente, L. and Solito, E. (2004) *Inflamm Res* 53, 125-32.
14. Lim, L.H. and Pervaiz, S. (2007) *FASEB J* 21, 968-75.

### Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

### Western Blot Buffer

**IMPORTANT:** For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween@ 20 at 4°C with gentle shaking, overnight.

### Applications Key

**W:** Western Blotting

### Cross-Reactivity Key

**H:** Human **M:** Mouse **R:** Rat **Mk:** Monkey

### Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

The Annexin V-FITC conjugate is a product manufactured for Cell Signaling Technology, Inc. by TAU Technologies BV.

All other trademarks are the property of their respective owners. Visit [cellsignal.com/trademarks](http://cellsignal.com/trademarks) for more information.

## Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.