Phospho-ALK (Tyr1604) Antibody



Orders: 877-616-CELL (2355)

orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com

cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Reactivity: H	Sensitivity: Endogenous	MW (kDa): 80 (NPM-ALK) 220 (ALK)	Source/Isotype: Rabbit	UniProt ID: #Q9UM73	Entrez-Gene Id: 238
	Application Western Blotting Simple Western™ Immunoprecipitatio	n		Dilution 1:1000 1:10 - 1:50 1:50	
			150 mM NaCl, 100 μg/	/ml BSA and 50% gl	ycerol. Store at –
sitivity					
ation	Polyclonal antibodies are produced by immunizing animals with a synthetic phosphopeptide corresponding to residues surrounding Tyr1604 of human ALK. Antibodies are purified by protein A and peptide affinity chromatography				
	Anaplastic lymphoma kinase (ALK) is a tyrosine kinase receptor for pleiotrophin (PTN), a growth factor involved in embryonic brain development (1-3). In ALK-expressing cells, PTN induces phosphorylation of both ALK and the downstream effectors IRS-1, Shc, PLCy, and PI3 kinase (1). ALK was originally discovered as a nucleophosmin (NPM)-ALK fusion protein produced by a translocation (4). Investigators have found that the NPM-ALK fusion protein is a constitutively active, oncogenic tyrosine kinase associated with anaplastic lymphoma (4). Research literature suggests that activation of PLCy by NPM-ALK may be a crucial step for its mitogenic activity and involved in the pathogenesis of anaplastic lymphomas (5). A distinct ALK oncogenic fusion protein involving ALK and echinoderm microtubule-associated protein like 4 (EML4) has been described in the research literature from a non-small cell lung cancer (NSCLC) cell line, with corresponding fusion transcripts present in some cases of lung adenocarcinoma. The short, amino-terminal region of the microtubule-associated protein EML4 is fused to the kinase domain of ALK (6-8). Phosphorylated Tyr664 of NPM-ALK (equivalent to Tyr1604 of full length ALK) is required for the interaction with PLCgamma (5). Site-directed mutagenesis of this tyrosine residue results in the loss of oncogenic activity of NPM-ALK (5).				
	of both ALK and the discovered as a nucl have found that the associated with anal ALK may be a crucia lymphomas (5). A distinct ALK oncog like 4 (EML4) has becell line, with corresshort, amino-termin of ALK (6-8). Phosphorylated Tyreinteraction with PLC	downstream effectors eophosmin (NPM)-ALK NPM-ALK fusion prote plastic lymphoma (4). Fill step for its mitogenic genic fusion protein inven described in the responding fusion transcrial region of the microt 664 of NPM-ALK (equiv. gamma (5). Site-directed	IRS-1, Shc, PLCy, and F fusion protein productin is a constitutively ac Research literature sug activity and involved in rolving ALK and echino earch literature from a ipts present in some coubule-associated protein alent to Tyr1604 of full	PI3 kinase (1). ALK ved by a translocation tive, oncogenic tyrogests that activation the pathogenesis derm microtubuleanon-small cell lungases of lung adenoiein EML4 is fused to length ALK) is required by a translocation to the second secon	vas originally on (4). Investigators osine kinase n of PLCy by NPM-of anaplastic associated protein g cancer (NSCLC) carcinoma. The o the kinase domain wired for the
		Application Western Blotting Simple Western™ Immunoprecipitatio Supplied in 10 mM s 20°C. Do not aliquot Sitivity Phospho-ALK (Tyr16 Tyr664 of NPM-ALK) including EGFR. Polyclonal antibodie corresponding to re and peptide affinity Anaplastic lymphom	Application Western Blotting Simple Western™ Immunoprecipitation Supplied in 10 mM sodium HEPES (pH 7.5), 20°C. Do not aliquot the antibody. Sitivity Phospho-ALK (Tyr1604) Antibody detects A Tyr664 of NPM-ALK). This antibody may creincluding EGFR. Polyclonal antibodies are produced by imm corresponding to residues surrounding Tyrand peptide affinity chromatography Anaplastic lymphoma kinase (ALK) is a tyro	H Endogenous 80 (NPM-ALK) 220 Rabbit Application Western Blotting Simple Western™ Immunoprecipitation Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 μg, 20°C. Do not aliquot the antibody. Sitivity Phospho-ALK (Tyr1604) Antibody detects ALK only when phospho Tyr664 of NPM-ALK). This antibody may cross-react with other act including EGFR. Polyclonal antibodies are produced by immunizing animals with a corresponding to residues surrounding Tyr1604 of human ALK. A and peptide affinity chromatography Anaplastic lymphoma kinase (ALK) is a tyrosine kinase receptor for	H Endogenous 80 (NPM-ALK) 220 Rabbit #Q9UM73 Application Dilution Western Blotting 1:1000 Simple Western™ 1:10 - 1:50 Immunoprecipitation 1:50 Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 μg/ml BSA and 50% gl 20°C. Do not aliquot the antibody. Sitivity Phospho-ALK (Tyr1604) Antibody detects ALK only when phosphorylated at Tyr1604 Tyr664 of NPM-ALK). This antibody may cross-react with other activated protein tyro including EGFR. Polyclonal antibodies are produced by immunizing animals with a synthetic phosphorylated affinity chromatography Anaplastic lymphoma kinase (ALK) is a tyrosine kinase receptor for pleiotrophin (PTN)

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

Applications Key

W: Western Blotting **W-S:** Simple Western[™] **IP:** Immunoprecipitation

Cross-Reactivity Key

H: Human

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.