

Store at
-20C
#33483**BOB-1/OBF-1 Antibody**

Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications:	Reactivity:	Sensitivity:	MW (kDa):	Source/Isotype:	UniProt ID:	Entrez-Gene Id:
W	H M R	Endogenous	35	Rabbit	#Q16633	5450

Product Usage Information**Application**

Western Blotting

Dilution

1:1000

Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA and 50% glycerol. Store at -20°C. Do not aliquot the antibody.

Specificity/Sensitivity

BOB-1/OBF-1 Antibody recognizes endogenous levels of total BOB-1/OBF-1 protein.

Species predicted to react based on 100% sequence homology

Dog, Pig

Source / Purification

Polyclonal antibodies are produced by immunizing animals with a synthetic peptide corresponding to residues surrounding Pro209 of human BOB-1/OBF-1 protein. Antibodies are purified by protein A and peptide affinity chromatography.

Background

B cell Oct binding factor-1 (BOB-1/OBF-1) is a B cell restricted transcriptional coactivator. BOB-1 facilitates transactivation of immunoglobulins and other B cell specific genes through the binding and activation of the transcription factors Oct-1 and Oct-2 (1-4). Research studies have demonstrated that BOB-1 expression is required for antigen-dependent B cell maturation (5-7). In pathological conditions such as classical Hodgkin disease, loss of BOB-1 expression is thought, in part, to contribute to the defect in immunoglobulin gene expression by Hodgkin and Reed Sternberg cells (8,9). In the context of multiple myeloma, overexpression of BOB-1 has been shown to contribute to malignant plasma cell growth, in part, through enhanced transactivation of *TNFRSF17/BCMA* (10).

Background References

1. Strubin, M. et al. (1995) *Cell* 80, 497-506.
2. Laumen, H. et al. (2000) *Eur J Immunol* 30, 458-69.
3. Gstaiger, M. et al. (1995) *Nature* 373, 360-2.
4. Brunner, C. and Wirth, T. (2006) *Nucleic Acids Res* 34, 1807-15.
5. Hess, J. et al. (2001) *Mol Cell Biol* 21, 1531-9.
6. Kim, U. et al. (1996) *Nature* 383, 542-7.
7. Schubart, D.B. et al. (1996) *Nature* 383, 538-42.
8. Re, D. et al. (2001) *Cancer Res* 61, 2080-4.
9. Stein, H. et al. (2001) *Blood* 97, 496-501.
10. Zhao, C. et al. (2008) *Oncogene* 27, 63-75.

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

Applications Key

W: Western Blotting

Cross-Reactivity Key

H: Human **M:** Mouse **R:** Rat

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless

separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.