#3360 Store at -20C

RagC Antibody



Orders:	877-616-CELL (2355) orders@cellsignal.com
Support:	877-678-TECH (8324)
Web:	info@cellsignal.com cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications: W, IP, IF-IC	Reactivity: H M R Mk	Sensitivity: Endogenous	MW (kDa): 50	Source/Isotype: Rabbit	UniProt ID: #Q9HB90	Entrez-Gene Id: 64121		
Product Usage Information		Application Western Blotting Immunoprecipitation Immunofluorescence	(Immunocytochem	istry)		Dilution 1:1000 1:50 1:400		
Storage		Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 μg/ml BSA and 50% glycerol. Store at – 20°C. Do not aliquot the antibody.						
Specificity/Sen	sitivity	RagC Antibody detects endogenous levels of total RagC protein.						
Source / Purification Polyclonal antibodies are produced by immunizing animals with a synthetic peptide residues near the amino-terminus of human RagC protein. Antibodies are purified peptide affinity chromatography.								
Background		The mTORC1 kinase complex is a critical regulator of cell growth (1,2). Its activity is modulated by energy levels, growth factors, and amino acids via signaling through Akt, MAPK, and AMPK pathways (3,4). Recent studies found that the four related GTPases, RagA, RagB, RagC, and RagD, interact with raptor within the mTORC1 complex (1,2). These interactions are both necessary and sufficient for mTORC1 activation in response to amino acid signals (1,2).						
Background Re	eferences	1. Sancak, Y. et al. (2008) <i>Science</i> 320, 1496-501. 2. Kim, E. et al. (2008) <i>Nat Cell Biol</i> 10, 935-45. 3. Hay, N. and Sonenberg, N. (2004) <i>Genes Dev</i> 18, 1926-45. 4. Wullschleger, S. et al. (2006) <i>Cell</i> 124, 471-84.						
Species Reactiv	vity	Species reactivity is determined by testing in at least one approved application (e.g., western blot).						
Western Blot E	Buffer	IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.						
Applications K	ey	W: Western Blotting IP: Immunoprecipitation IF-IC: Immunofluorescence (Immunocytochemistry)						
Cross-Reactivi	ty Key	H: Human M: Mouse R: Rat Mk: Monkey						
Trademarks ar	nd Patents	Cell Signaling Technol	hnology is a trademark of Cell Signaling Technology, Inc.					
		All other trademarks a more information.	are the property of t	heir respective owners.	Visit cellsignal.com	n/trademarks for		
Limited Uses		Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.						
		approved, cleared, or purpose. Customer sh any manner that conf Customer as the end- diagnostic, prophylact	licensed by the FDA nall not use any Proo licts with its labeling user and solely for i tic or therapeutic pu	se Only or a similar labe or other regulatory fore duct for any diagnostic o statement. Products so research and developme urposes, or any purchase e, requires a separate lic	eign or domestic en or therapeutic purp old or licensed by C ent uses. Any use o e of Product for res	ntity, for any ose, or otherwise in ST are provided for f Product for ale (alone or as a		

sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.