

Phospho-eIF2 α (Ser51) (D9G8) XPTM Rabbit mAb

Orders ■ 877-616-CELL (2355)
orders@cellsignal.com

Support ■ 877-678-TECH (8324)
info@cellsignal.com

Web ■ www.cellsignal.com

rev. 01/11/16

For Research Use Only. Not For Use In Diagnostic Procedures.

Applications	Species Cross-Reactivity*	Molecular Wt.	Isotype
W, IP, IHC-P Endogenous	H, M, R, Mk, Dm	38 kDa	Rabbit IgG**


Background: Phosphorylation of the eukaryotic initiation factor 2 (eIF2) α subunit is a well-documented mechanism to downregulate protein synthesis under a variety of stress conditions. Eukaryotic initiation factor 2 binds GTP and Met-tRNAi and transfers Met-tRNA to the 40S subunit to form the 43S preinitiation complex (1,2). eIF2 promotes a new round of translation initiation by exchanging GDP for GTP, a reaction catalyzed by eIF2B (1,2). Kinases that are activated by viral infection (PKR), endoplasmic reticulum stress (PERK/PEK), amino acid deprivation (GCN2) or heme deficiency (HRI) can phosphorylate the α subunit of eIF2 (3,4). This phosphorylation stabilizes the eIF2-GDP-eIF2B complex and inhibits the turnover of eIF2B. Induction of PKR by IFN- γ and TNF- α induces potent phosphorylation of eIF2 α at Ser51 (5,6).

Specificity/Sensitivity: Phospho-eIF2 α (Ser51) (D9G8) XPTM Rabbit mAb detects endogenous eIF2 α only when phosphorylated at Ser51. The antibody does not recognize eIF2 α phosphorylated at other sites.


Source/Purification: Monoclonal antibody is produced by immunizing animals with a synthetic phosphopeptide corresponding to residues surrounding Ser51 of human eIF2 α .

Background References:

- (1) Kimball, S.R. (1999) *Int. J. Biochem. Cell Biol.* 31, 25–29.
- (2) De Haro, C. et al. (1996) *FASEB J.* 10, 1378–1387.
- (3) Kaufman, R.J. (1999) *Genes Dev.* 13, 1211–1233.
- (4) Sheikh, M.S. and Fornace Jr., A.J. (1999) *Oncogene* 18, 6121–6128.
- (5) Cheshire, J.L. et al. (1999) *J. Biol. Chem.* 274, 4801–4806.
- (6) Zamanian-Daryoush, M. et al. (2000) *Mol. Cell. Biol.* 20, 1278–1290.



Western blot analysis of extracts from C2C12 cells, untreated or thapsigargin-treated, using Phospho-eIF2 α (Ser51) (D9G8) XPTM Rabbit mAb (upper) or eIF2 α Antibody #9722 (lower).



Immunohistochemical analysis of paraffin-embedded human lymphoma using Phospho-eIF2 α (Ser51) (D9G8) XPTM Rabbit mAb.

Entrez-Gene ID #1965
Swiss-Prot Acc. #P05198

Storage: Supplied in 10mM HEPES (pH 7.5), 150 mM NaCl, 100 μ g/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at -20°C. Do not aliquot the antibody.

*Species cross-reactivity is determined by western blot.

**Anti-rabbit secondary antibodies must be used to detect this antibody.



Recommended Antibody Dilutions:

Western blotting	1:1000
Immunoprecipitation	1:100
Immunohistochemistry (Paraffin)	1:100
Unmasking buffer:	Citrate
Antibody diluent:	SignalStain [®] Antibody Diluent #8112


For application specific protocols please see the web page for this product at www.cellsignal.com.

Please visit www.cellsignal.com for a complete listing of recommended companion products.

IMPORTANT: For western blots, incubate membrane with diluted antibody in 5% w/v BSA, 1X TBS, 0.1% Tween-20 at 4°C with gentle shaking, overnight.



Immunohistochemical analysis of paraffin-embedded human colon carcinoma, untreated (left) or λ phosphatase-treated (right), using Phospho-eIF2 α (Ser51) (D9G8) XP™ Rabbit mAb.



Immunohistochemical analysis of paraffin-embedded human lung carcinoma using Phospho-eIF2 α (Ser51) (D9G8) XP™ Rabbit mAb.