

Atg3 Antibody



Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications:	Reactivity:	Sensitivity:	MW (kDa):	Source/Isotype:	UniProt ID:	Entrez-Gene Id:
W	H M R Mk	Endogenous	40	Rabbit	#Q9NT62	64422

Product Usage Information

Application

Western Blotting

Dilution

1:1000

Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA and 50% glycerol. Store at -20°C. Do not aliquot the antibody.

Specificity/Sensitivity

Atg3 Antibody detects endogenous levels of total Atg3 protein.

Species predicted to react based on 100% sequence homology

Chicken, Xenopus, Bovine, Dog

Source / Purification

Polyclonal antibodies are produced by immunizing animals with a synthetic peptide corresponding to residues near the amino terminus of Atg3. Antibodies are purified by protein A and peptide affinity chromatography.

Background

Autophagy is a catabolic process for the autophagosomic-lysosomal degradation of bulk cytoplasmic contents (1). The molecular machinery of autophagy was largely discovered in yeast and referred to as autophagy-related genes (Atg). Formation of the autophagic vesicles involves two ubiquitin-like conjugation systems, Atg12-Atg5 and Atg8-phosphatidylethanolamine (Atg8-PE), which are essential for autophagy and widely conserved in eukaryotes (2). There are at least three Atg8 homologs in mammalian cells, GATE-16, GABARAP, and LC3, that are conjugated by lipids (3,4). Lipid conjugation of Atg8 and its mammalian homologs requires Atg3 (Apg3p/Aut1p in yeast), an ubiquitously expressed E2-like enzyme (5-7). Following C-terminal cleavage by the cysteine protease Atg4, the exposed glycine residue of Atg8 binds to the E1-like enzyme Atg7, is transferred to Atg3, and then conjugated to phosphatidylethanolamine. Atg3-deficient mice die within 1 day after birth and are completely defective for the conjugation of Atg8 homologs and autophagosome formation (8).

Background References

1. Reggiori, F. and Klionsky, D.J. (2002) *Eukaryot Cell* 1, 11-21.
2. Ohsumi, Y. (2001) *Nat Rev Mol Cell Biol* 2, 211-6.
3. Kabeya, Y. et al. (2000) *EMBO J* 19, 5720-8.
4. Kabeya, Y. et al. (2004) *J Cell Sci* 117, 2805-12.
5. Tanida, I. et al. (2002) *J Biol Chem* 277, 13739-44.
6. Ichimura, Y. et al. (2000) *Nature* 408, 488-92.
7. Schlumpberger, M. et al. (1997) *J Bacteriol* 179, 1068-76.
8. Sou, Y.S. et al. (2008) *Mol Biol Cell* 19, 4762-75.

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

Applications Key

W: Western Blotting

Cross-Reactivity Key

H: Human **M:** Mouse **R:** Rat **Mk:** Monkey

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's

terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.