RECK (D8C7) Rabbit mAb	C T	Cell Signaling	
	Orders:	877-616-CELL (2355) orders@cellsignal.com	
	Support:	877-678-TECH (8324)	
	Web:	info@cellsignal.com cellsignal.com	



3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications: W	<b>Reactivity:</b> H M R Mk	<b>Sensitivity:</b> Endogenous	<b>MW (kDa):</b> 110	<b>Source/Isotype:</b> Rabbit IgG	<b>UniProt ID:</b> #O95980	Entrez-Gene Id: 8434		
Product Usage Information		<b>Application</b> Western Blotting			Dilution 1:1000			
Storage		Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 μg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at –20°C. Do not aliquot the antibody.						
Specificity/Sen	sitivity	RECK (D8C7) Rabbit mAb detects endogenous levels of total RECK protein.						
Source / Purifi	cation	Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to residues surrounding Ile913 of human RECK protein.						
Background		RECK (reversion-inducing cysteine-rich protein with Kazal motif) is a GPI-anchored membrane glycoprotein that negatively regulates members of the matrix metalloproteinase (MMP) family and functions as a suppressor of transformation (1,2). Its function in MMP inhibition makes RECK a crucial factor in the regulation of extracellular matrix formation and stability during development (2-4). RECK has also been linked to the regulation of other extracellar matrix proteases such as ADAM10 and CD13 and functions in modulating target protein endocytosis and Notch signaling (5,6). RECK is widely expressed in normal tissue and decreased expression of RECK due to promoter methylation has been correlated with tumor transformation, angiogenesis and metastasis (1,7-9). Therefore, loss of RECK expression serves as a prognostic hallmark for cancer malignancy (10,11)						
Background R	eferences	1. Takahashi, C. et al. ( 2. Oh, J. et al. (2001) <i>C</i> 3. Kondo, S. et al. (200 4. Echizenya, M. et al. 5. Miki, T. et al. (2007) 6. Muraguchi, T. et al. 7. Noda, M. et al. <i>Can</i> 8. Long, N.K. et al. (20 9. Chang, H.C. et al. (20 11. Noda, M. and Taka	<i>ell</i> 107, 789-800. )7) <i>J Cell Sci</i> 120, 84 (2005) <i>Oncogene 2 J Biol Chem</i> 282, 12 (2007) <i>Nat Neuros cer Metastasis Rev</i> 08) <i>Oral Oncol</i> 44, 007) <i>Cancer Sci</i> 98, 07) <i>Cancer Metasta</i>	4, 5850-7. 2341-52. ci 10, 838-45. 22, 167-75. 1052-8. 169-73. asis Rev 26, 675-83.				
Species Reacti	vity	Species reactivity is de	etermined by testir	g in at least one approve	ed application (e.g.,	western blot).		
Western Blot E	Buffer	IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.						
Applications K	ey	W: Western Blotting						
Cross-Reactivi	ty Key	H: Human M: Mouse R: Rat Mk: Monkey						
Trademarks ar	nd Patents	Cell Signaling Technol	ogy is a trademark	of Cell Signaling Techno	logy, Inc.			
		All other trademarks a more information.	are the property of	their respective owners.	Visit cellsignal.com	ı/trademarks for		
Limited Uses		Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.						

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.