

Store at
-20C
#36067**RanGAP1 (D2T7T) Rabbit mAb**

Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications: W, IP, eCLIP	Reactivity: H M R Mk	Sensitivity: Endogenous	MW (kDa): 68, 82	Source/Isotype: Rabbit IgG	UniProt ID: #P46060	Entrez-Gene Id: 5905
--------------------------------------	--------------------------------	-----------------------------------	----------------------------	--------------------------------------	-------------------------------	--------------------------------

Product Usage Information**Application**

Western Blotting
Immunoprecipitation
eCLIP

Dilution

1:1000
1:100
1:200

For more information about the RBP-eCLIP service please visit Eclipsebio.

Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at -20°C. Do not aliquot the antibody.

Specificity/Sensitivity

RanGAP1 (D2T7T) Rabbit mAb recognizes endogenous levels of total RanGAP1 protein. This antibody detects unmodified and SUMO-1 modified RanGAP1 protein.

Source / Purification

Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to residues near the amino terminus of human RanGAP1 protein.

Background

Ran GTPase activating protein 1 (RanGAP1) regulates GTP hydrolysis by Ran, which is required for the ability of Ran to regulate nucleocytoplasmic shuttling (1). A significant fraction of cellular RanGAP1 is covalently modified by SUMO-1, which is required for relocalization of RanGAP1 from the cytoplasm to the nuclear pore complex and the mitotic spindle (2-5). Research studies demonstrate that RanGAP1 sumoylation is required for stable association of RanGAP1 with RanBP2. Together with the SUMO-conjugating enzyme UBC9/UBE2I, RanGAP1 and RanBP2 are part of a SUMO E3 ligase complex that is implicated in regulating nucleocytoplasmic protein trafficking (6-8). Phosphorylation of RanGAP1 occurs in a cell-cycle-dependent manner and may play a role in regulating RanGAP1 catalytic activity (9,10).

Background References

1. Bischoff, F.R. et al. (1994) *Proc Natl Acad Sci U S A* 91, 2587-91.
2. Mahajan, R. et al. (1997) *Cell* 88, 97-107.
3. Matunis, M.J. et al. (1998) *J Cell Biol* 140, 499-509.
4. Matunis, M.J. et al. (1996) *J Cell Biol* 135, 1457-70.
5. Joseph, J. et al. (2002) *J Cell Biol* 156, 595-602.
6. Hutten, S. et al. (2008) *Mol Biol Cell* 19, 2300-10.
7. Werner, A. et al. (2012) *Mol Cell* 46, 287-98.
8. Zhu, S. et al. (2009) *Mol Cell* 33, 570-80.
9. Swaminathan, S. et al. (2004) *J Cell Biol* 164, 965-71.
10. Takeda, E. et al. (2005) *Cell Struct Funct* 30, 69-80.

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

Applications Key

W: Western Blotting **IP:** Immunoprecipitation **eCLIP:** eCLIP

Cross-Reactivity Key

H: Human **M:** Mouse **R:** Rat **Mk:** Monkey

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

XP is a registered trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.