

## 36233

## PSD95 (7E3) Mouse mAb



Orders: 877-616-CELL (2355)

orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com

cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

## For Research Use Only. Not for Use in Diagnostic Procedures.

<b>Applications:</b> W, IF-F, IF-IC	Reactivity: H M R	<b>Sensitivity:</b> Endogenous	<b>MW (kDa):</b> 95	<b>Source/Isotype:</b> Mouse IgG1	<b>UniProt ID:</b> #P78352	Entrez-Gene Id: 1742
Product Usage Information		Application Western Blotting Immunofluorescence	` '			<b>Dilution</b> 1:1000 1:100
Storage		Immunofluorescence (Immunocytochemistry)  Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at –20°C. Do not aliquot the antibody.  For a carrier-free (BSA and azide free) version of this product see product #37350.				
Specificity/Sensitivity		PSD95 (7E3) Mouse mAb recognizes endogenous levels of total PSD95 protein.				
Source / Purification		Monoclonal antibody is produced by immunizing animals with recombinant protein specific to the full length rat PSD95 protein.				
Background		Postsynaptic Density protein 95 (PSD95) is a member of the membrane-associated guanylate kinase (MAGUK) family of proteins. These family members consist of an amino-terminal variable segment followed by three PDZ domains, an SH3 domain, and an inactive guanylate kinase (GK) domain. PSD95 is a scaffolding protein involved in the assembly and function of the postsynaptic density complex (1-2). PSD95 participates in synaptic targeting of AMPA receptors through an indirect manner involving stargazin and related transmembrane AMPA receptor regulatory proteins (TARPs) (3). It is implicated in experience-dependent plasticity and plays an indispensable role in learning (4). Mutations in PSD95 are associated with autism (5).				
Background R	eferences	1. Cao, J. et al. (2005) <i>J. Cell Biol</i> 168, 117-26. 2. Chetkovich, D.M. et al. (2002) <i>J. Neurosci.</i> 22, 6415-25. 3. Cai, C. et al. (2006) <i>J. Biol. Chem.</i> 281, 4267-73. 4. Yao, W.D. et al. (2004) <i>Neuron</i> 41, 625-38. 5. Cline, H. (2005) <i>Curr. Biol.</i> 15, R203-5.				
Species Reacti	vity	Species reactivity is d	etermined by testin	g in at least one approve	ed application (e.g.,	western blot).

Species reactivity is determined by testing in at least one approved application (e.g., western biot)

Western Blot Buffer IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X

TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

**Applications Key** W: Western Blotting **IF-F**: Immunofluorescence (Frozen) **IF-IC**: Immunofluorescence

(Immunocytochemistry)

Cross-Reactivity Key H: Human M: Mouse R: Rat

**Trademarks and Patents** Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

XP is a registered trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for a constant of their respective owners are the property of their respective owners.

more information.

Limited Uses Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST,

the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless

separately accepted in writing by a legally authorized representative of CST, are rejected and are of no

force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.