



**Orders:** 877-616-CELL (2355)  
orders@cellsignal.com

**Support:** 877-678-TECH (8324)

**Web:** info@cellsignal.com  
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

Store at -20C  
#3664

## RhoE (4) Mouse mAb

**For Research Use Only. Not for Use in Diagnostic Procedures.**

<b>Applications:</b> W	<b>Reactivity:</b> H Mk	<b>Sensitivity:</b> Endogenous	<b>MW (kDa):</b> 29	<b>Source/Isotype:</b> Mouse IgG1	<b>UniProt ID:</b> #P61587	<b>Entrez-Gene Id:</b> 390
---------------------------	----------------------------	-----------------------------------	------------------------	--------------------------------------	-------------------------------	-------------------------------

### Product Usage Information

#### Application

Western Blotting

#### Dilution

1:1000

### Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at -20°C. Do not aliquot the antibody.

### Specificity/Sensitivity

RhoE (4) Mouse mAb detects endogenous levels of total RhoE protein.

### Source / Purification

Monoclonal antibody is produced by immunizing animals with full length recombinant human RhoE.

### Background

Rho family small GTPases act as molecular switches that regulate processes such as cell migration, adhesion, proliferation and differentiation. Typically, they are activated by guanine nucleotide exchange factors (GEFs), which catalyze the exchange of bound GDP for GTP, and are inhibited by GTPase activating proteins (GAPs), which catalyze the hydrolysis of GTP to GDP (1). Rnd1, Rnd2 and RhoE/Rnd3 comprise the evolutionarily divergent Rnd family of Rho-type small GTPases, which lack GTPase activity and therefore remain in a GTP-bound state (2, reviewed in 3). RhoE/Rnd3 activity leads to a decrease in stress fibers and increased cell migration, at least in part through regulation of the Rho-dependent kinase ROCK1 (4). Activity of RhoE/Rnd3 itself is regulated by ROCK1, which phosphorylates RhoE/Rnd3 at Ser11, enhancing its activity (5). RhoE/Rnd3 has been implicated in inhibition of DNA-damage induced apoptosis (6) and cell cycle arrest (7). In keratinocytes, RhoE/Rnd3 regulates differentiation through its effects on both proliferation and adhesion (8).

### Background References

1. DerMardirossian, C. and Bokoch, G.M. (2005) *Trends Cell Biol* 15, 356-63.
2. Foster, R. et al. (1996) *Mol Cell Biol* 16, 2689-99.
3. Riento, K. et al. (2005) *Biochem Soc Trans* 33, 649-51.
4. Riento, K. et al. (2003) *Mol Cell Biol* 23, 4219-29.
5. Riento, K. et al. (2005) *EMBO J* 24, 1170-80.
6. Boswell, S.A. et al. (2007) *J Biol Chem* 282, 4850-8.
7. Poch, E. et al. (2007) *Exp Cell Res* 313, 719-31.
8. Liebig, T. et al. (2009) *Mol Biol Cell* 20, 452-63.

### Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

### Western Blot Buffer

**IMPORTANT:** For western blots, incubate membrane with diluted primary antibody in 5% w/v nonfat dry milk, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

### Applications Key

**W:** Western Blotting

### Cross-Reactivity Key

**H:** Human **Mk:** Monkey

### Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit [cellsignal.com/trademarks](http://cellsignal.com/trademarks) for more information.

### Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any

purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.