

**Phospho-GRB10 (Tyr67) Antibody**

**Orders:** 877-616-CELL (2355)  
orders@cellsignal.com

**Support:** 877-678-TECH (8324)

**Web:** info@cellsignal.com  
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

**For Research Use Only. Not for Use in Diagnostic Procedures.**

Applications:	Reactivity:	Sensitivity:	MW (kDa):	Source/Isotype:	UniProt ID:	Entrez-Gene Id:
W	H	Transfected Only	76 (Transfected only)	Rabbit	#Q13322	2887

<b>Product Usage Information</b>	<b>Application</b> Western Blotting	<b>Dilution</b> 1:1000
<b>Storage</b>	Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA and 50% glycerol. Store at -20°C. Do not aliquot the antibody.	
<b>Specificity/Sensitivity</b>	Phospho-GRB10 (Tyr67) Antibody detects transfected GRB10 only when phosphorylated at tyrosine 67. The antibody does not cross-react with phosphorylated GRB7 or GRB14.	
<b>Source / Purification</b>	Polyclonal antibodies are produced by immunizing animals with a synthetic phosphopeptide corresponding to residues surrounding Tyr67 of human GRB10. Antibodies are purified by protein A and peptide affinity chromatography.	
<b>Background</b>	The GRB7 family of adaptor proteins consist of GRB7, GRB10 and GRB14, which all contain an amino-terminal proline-rich SH3 binding domain, followed by PH, PBS, and SH2 domains. Each member of the family has several splice variants (1). It has been reported that GRB10 interacts with many receptor tyrosine kinases (RTKs) as well as downstream signal molecules including Raf, Akt, and Nedd4 (1,2). Although it was originally thought that GRB10 is exclusively phosphorylated at serine residues (3), Src kinase family members have been shown to phosphorylate GRB10 at Tyr67 (4).	
<b>Background References</b>	1. Han, D. C. et al. (2001) <i>Oncogene</i> 20, 6315-6321. 2. Jahn, T. et al. (2002) <i>Mol. Cell. Biol.</i> 22, 979-991. 3. Ooi, J. et al. (1995) <i>Oncogene</i> 10, 1621-1630. 4. Langlais, P. et al. (2000) <i>Oncogene</i> 19, 2895-2903.	

<b>Species Reactivity</b>	Species reactivity is determined by testing in at least one approved application (e.g., western blot).
<b>Western Blot Buffer</b>	IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.
<b>Applications Key</b>	<b>W:</b> Western Blotting
<b>Cross-Reactivity Key</b>	<b>H:</b> Human
<b>Trademarks and Patents</b>	Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.  All other trademarks are the property of their respective owners. Visit <a href="http://cellsignal.com/trademarks">cellsignal.com/trademarks</a> for more information.
<b>Limited Uses</b>	<p>Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.</p> <p>Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a</p>

component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.