



Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

Store at -20C
#3719

GNB3 Antibody

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications: W	Reactivity: H M R	Sensitivity: Endogenous	MW (kDa): 31	Source/Isotype: Rabbit	UniProt ID: #P16520	Entrez-Gene Id: 2784
---------------------------	-----------------------------	-----------------------------------	------------------------	----------------------------------	-------------------------------	--------------------------------

Product Usage Information

Application

Western Blotting

Dilution

1:1000

Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA and 50% glycerol. Store at -20°C. Do not aliquot the antibody.

Specificity/Sensitivity

GNB3 Antibody recognizes endogenous levels of total GNB3 protein.

Source / Purification

Polyclonal antibodies are produced by immunizing animals with a synthetic peptide corresponding to residues surrounding Ile123 of human GNB3 protein. Antibodies are purified by protein A and peptide affinity chromatography.

Background

Heterotrimeric guanine nucleotide-binding proteins, G proteins, transduce ligand binding to G protein-coupled receptors (GPCRs) into intracellular responses (1). G proteins are comprised of 3 subunits, alpha (Gα), beta (Gβ), and gamma (Gγ). Upon activation of GPCRs, the receptor promotes the exchange of GDP to GTP of Gα, changing the confirmation of the switch regions within Gα. The receptor bound heterotrimeric G protein (inactive) is then released, and dissociates into the GTP-bound Gα (active) monomer and the Gβ/Gγ heterodimer (1,2). Gα activates adenylyl cyclase, which converts ATP to the second messenger cAMP. Gα also activates phosphoinositide-specific phospholipase C (PLC), which catalyzes hydrolysis of the phospholipid of phosphatidylinositol 4,5-bisphosphate (PIP₂), releasing the second messengers IP₃ and 1,2-diacylglycerol (DAG). IP₃ activates IP₃ receptors to release Ca²⁺ from the ER. DAG is an activator of protein kinase C (PKC), which in turn activates the Erk1/2 pathway (1,3). The primary function of the Gβ/Gγ heterodimer is to inhibit Gα, although it may also activate second messengers (e.g. PLC pathway) or gate ion channels (e.g. GIRK) (1). Guanine nucleotide-binding protein b3 (GNB3) is an isoform of the b subunit. Research studies have shown that a polymorphism in the GNB3 gene, C825T, is associated with hypertension, obesity, and depression (4).

Background References

- Hamm, H.E. (1998) *J Biol Chem* 273, 669-72.
- Ritchey, E.R. et al. (2010) *Neuroscience* 169, 1376-91.
- Hisatsune, C. et al. (2005) *J Biol Chem* 280, 11723-30.
- Roskopf, D. et al. (2000) *Hypertension* 36, 33-41.

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween@ 20 at 4°C with gentle shaking, overnight.

Applications Key

W: Western Blotting

Cross-Reactivity Key

H: Human **M:** Mouse **R:** Rat

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.