

#3775 Store at -20C

Jak3 Antibody



Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

| | | | | | | |
|---------------------------|-------------------------|-----------------------------------|-------------------------|----------------------------------|-------------------------------|--------------------------------|
| Applications: W | Reactivity: H | Sensitivity: Endogenous | MW (kDa): 115 | Source/Isotype: Rabbit | UniProt ID: #P52333 | Entrez-Gene Id: 3718 |
|---------------------------|-------------------------|-----------------------------------|-------------------------|----------------------------------|-------------------------------|--------------------------------|

| | | |
|----------------------------------|--|---------------------------|
| Product Usage Information | Application Western Blotting | Dilution 1:1000 |
| Storage | Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA and 50% glycerol. Store at -20°C. Do not aliquot the antibody. | |
| Specificity/Sensitivity | Jak3 Antibody detects endogenous levels of total Jak3 protein. No cross-reactivity was detected with other family members at physiological conditions. | |
| Source / Purification | Polyclonal antibodies are produced by immunizing animals with a synthetic peptide corresponding to residues at the carboxy-terminus of Jak3. Antibodies are purified by protein A and peptide affinity chromatography. | |
| Background | Members of the Janus family of tyrosine kinases (Jak1, Jak2, Jak3, and Tyk2) are activated by ligands binding to a number of associated cytokine receptors (1). Upon cytokine receptor activation, Jak proteins become autophosphorylated and phosphorylate their associated receptors to provide multiple binding sites for signaling proteins. These associated signaling proteins, such as Stats (2), Shc (3), insulin receptor substrates (4), and focal adhesion kinase (FAK) (5), typically contain SH2 or other phospho-tyrosine-binding domains. | |
| Background References | <ol style="list-style-type: none"> 1. Leonard, W.J. and O'Shea, J.J. (1998) <i>Annu Rev Immunol</i> 16, 293-322. 2. Darnell, J.E. (1997) <i>Science</i> 277, 1630-5. 3. VanderKuur, J. et al. (1995) <i>J Biol Chem</i> 270, 7587-93. 4. Argetsinger, L.S. et al. (1995) <i>J Biol Chem</i> 270, 14685-92. 5. Zhu, T. et al. (1998) <i>J Biol Chem</i> 273, 10682-9. | |

| | |
|-------------------------------|---|
| Species Reactivity | Species reactivity is determined by testing in at least one approved application (e.g., western blot). |
| Western Blot Buffer | IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight. |
| Applications Key | W: Western Blotting |
| Cross-Reactivity Key | H: Human |
| Trademarks and Patents | Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc. All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information. |
| Limited Uses | <p>Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.</p> <p>Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not</p> |

sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.