

SNF2H (D4W6N) Rabbit mAb

Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications: W	Reactivity: H	Sensitivity: Endogenous	MW (kDa): 125	Source/Isotype: Rabbit IgG	UniProt ID: #O60264	Entrez-Gene Id: 8467
---------------------------	-------------------------	-----------------------------------	-------------------------	--------------------------------------	-------------------------------	--------------------------------

Product Usage Information**Application**

Western Blotting

Dilution

1:1000

Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at -20°C. Do not aliquot the antibody.

Specificity/Sensitivity

SNF2H (D4W6N) Rabbit mAb recognizes endogenous levels of total SNF2H protein.

Source / Purification

Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to residues surrounding Ala65 of human SNF2H protein.

Background

Sucrose nonfermenting 2 homolog (SNF2H, SMARCA5) is one of two orthologs of the ISWI (imitation switch) ATPases encoded by the mammalian genome (1). SNF2H is part of the SNF2 family of chromatin remodeling factors that use ATP hydrolysis to catalyze biochemical reactions in several mammalian chromatin-remodeling complexes, including ACF1, RSF1, CHRAC, NoRC, WSTF, and WCRF180 (2). Research studies show that SNF2H is crucial for chromatin organization, DNA damage response, and differentiation (1-7). The SNF2H helicase facilitates DNA damage repair by actively moving nucleosomes for DNA damage response (DDR) proteins to effectively associate with damaged regions (3). Additional studies show that repair of double stranded breaks (DSBs) significantly decreases in the absence of SNF2H (3), and these cells become highly sensitive to DNA damage caused by x-rays and chemical treatments inducing DSBs (4,5).

Background References

1. Lazzaro, M.A. and Picketts, D.J. (2001) *J Neurochem* 77, 1145-56.
2. Kasten, M.M. et al. (2011) *Cell* 144, 310.e1.
3. Helfricht, A. et al. (2013) *Cell Cycle* 12, 3070-82.
4. Mueller, A.C. et al. (2013) *Oncogene* 32, 1164-72.
5. Lan, L. et al. (2010) *Mol Cell* 40, 976-87.
6. Smeenk, G. et al. (2013) *J Cell Sci* 126, 889-903.
7. Chioda, M. et al. (2010) *Development* 137, 3513-22.

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

Applications Key

W: Western Blotting

Cross-Reactivity Key

H: Human

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

CST is a registered trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.