

PARN (A42) Antibody



Orders: 877-616-CELL (2355)

orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com

cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications: W	Reactivity: H M R Mk	Sensitivity: Endogenous	MW (kDa): 78	Source/Isotype: Rabbit	UniProt ID: #O95453	Entrez-Gene Id: 5073
Product Usage Information		Application Western Blotting			Dilution 1:1000	
Storage		Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 μg/ml BSA and 50% glycerol. Store at – 20°C. Do not aliquot the antibody.				
Specificity/Sensitivity		PARN (A42) Antibody detects endogenous levels of total PARN protein.				
Source / Purification		Polyclonal antibodies are produced by immunizing animals with a synthetic peptide corresponding to amino acid sequence surrounding Ala42 of human PARN. Antibodies are purified by Protein A and peptide affinity chromatography.				
Background		Cellular levels of mRNAs are controlled by mRNA stability, the rate of synthesis and the rate of degradation. The presence and length of the poly(A) tail has been associated with mRNA stability (1). Exonucleolytic shortening of the poly(A) tail is the process that initiates the decay of many eukaryotic mRNAs (2). Poly(A)-specific ribonuclease (PARN) is the enzyme responsible for initiation of deadenylation and exonucleolytic shortening of mRNA transcripts. Through an evolutionarily conserved mechanism, PARN also translationally silences selective mRNAs during early embryonic development (3). PARN is constitutively expressed in most mammalian tissues and plays a critical role in the post-transcriptional control of gene expression (4).				
Background References		1. Morales, J. et al. (1997) <i>J Biol Chem</i> 272, 6607-13. 2. Wilson, T. and Treisman, R. (1988) <i>Nature</i> 336, 396-9. 3. Sachs, A.B. et al. (1997) <i>Cell</i> 89, 831-8. 4. Fritz, D.T. et al. (2004) <i>Cell Biochem Biophys</i> 41, 265-78.				
Species Reactivity		Species reactivity is determined by testing in at least one approved application (e.g., western blot).				
Western Blot Buffer		IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.				
Applications Key		W: Western Blotting				
Cross-Reactivity Key		H: Human M: Mouse R: Rat Mk: Monkey				
Trademarks and Patents		Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.				
		All other trademarks a more information.	re the property of t	heir respective owners.	Visit cellsignal.com	/trademarks for
Limited Uses		Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.				
				se Only or a similar labe or other regulatory fore		

purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not

sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.