e at -20C	MBD3 Antibody		ell Signaling
Store		Orders:	877-616-CELL (2355) orders@cellsignal.com
		Support:	877-678-TECH (8324)
896		Web:	info@cellsignal.com cellsignal.com
#3		3 Trask Lane Danvers Mass	achusetts 01923 USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications: W	Reactivity: H M R Mk	Sensitivity: Endogenous	MW (kDa): 28, 32, 34	Source/Isotype: Rabbit	UniProt ID: #O95983	Entrez-Gene Id: 53615	
Product Usage Information		Application Western Blotting			Dilution 1:1000		
Storage		Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 μg/ml BSA and 50% glycerol. Store at – 20°C. Do not aliquot the antibody.					
Specificity/Sensitivity		MBD3 Antibody detects endogenous levels of all three isoforms of the MBD3 protein (MBD3A, MBD3B, MBD3C). This antibody does not cross react with MBD2.					
Source / Purification		Polyclonal antibodies are produced by immunizing animals with a synthetic peptide corresponding to the human MBD3 protein. Antibodies are purified by protein A and peptide affinity chromatography.					
Background		Methyl-CpG-binding protein 2 (MeCP2) is the founding member of a family of methyl-CpG-binding domain (MBD) proteins that also includes MBD1, MBD2, MBD3, MBD4, MBD5, and MBD6 (1-3). Apart from MBD3, these proteins bind methylated cytosine residues in the context of the di-nucleotide 5'-CG-3' to establish and maintain regions of transcriptionally inactive chromatin by recruiting a variety of co-repressor proteins (2). MeCP2 recruits histone deacetylases HDAC1 and HDAC2, and the DNA methyltransferase DNMT1 (4-6). MBD1 couples transcriptional silencing to DNA replication and interacts with the histone methyltransferases ESET and SUV39H1 (7,8). MBD2 and MBD3 co-purify as part of the NuRD (nucleosome remodeling and histone de-acetylation) co-repressor complex, which contains the chromatin remodeling ATPase Mi-2, HDAC1, and HDAC2 (9,10). MBD5 and MBD6 have recently been identified and little is known regarding their protein interactions. MBD proteins are associated with cancer and other diseases; MBD4 is best characterized for its role in DNA repair and MBD2 has been linked to intestinal cancer (11,12). Mutations in the <i>MeCP2</i> gene cause the neurologic developmental disorder Rett Syndrome (13). MeCP2 protein levels are high in neurons, where it plays a critical role in multiple synaptic processes (14). In response to various physiological stimuli, MeCP2 is phosphorylated on Ser421 and regulates the expression of genes controlling dendritic patterning and spine morphogenesis (14). Disruption of this process in individuals with altered MeCP2 may cause the pathological changes seen in Rett Syndrome.					
Background References		 Clouaire, T. and Stancheva, I. (2008) <i>Cell Mol Life Sci</i> 65, 1509-22. Hendrich, B. and Bird, A. (1998) <i>Mol Cell Biol</i> 18, 6538-47. Roloff, T.C. et al. (2003) <i>BMC Genomics</i> 4, 1. Nan, X. et al. (1998) <i>Nature</i> 393, 386-9. Jones, P.L. et al. (1998) <i>Nat Genet</i> 19, 187-91. Fuks, F. et al. (2003) <i>J Biol Chem</i> 278, 4035-40. Sarraf, S.A. and Stancheva, I. (2004) <i>Mol Cell</i> 15, 595-605. Fujita, N. et al. (2003) <i>J Biol Chem</i> 278, 24132-8. Zhang, Y. et al. (1999) <i>Genes Dev</i> 13, 1924-35. Wade, P.A. et al. (1999) <i>Nat Genet</i> 23, 62-6. Hendrich, B. et al. (1999) <i>Nat Genet</i> 34, 145-7. Sansom, O.J. et al. (2003) <i>Nat Genet</i> 34, 145-7. Miltenberger-Miltenyi, G. and Laccone, F. (2003) <i>Hum Mutat</i> 22, 107-15. Zhou, Z. et al. (2006) <i>Neuron</i> 52, 255-69. 					
Species Reacti	ivity	Species reactivity is de	etermined by testin	g in at least one approve	ed application (e.g.,	western blot).	
Western Blot Buffer		IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.					
Applications K	(ey	W: Western Blotting					
Cross-Reactivi	ity Key	H: Human M: Mouse	R: Rat Mk: Monkey				

Trademarks and Patents	Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc. All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.
Limited Uses	Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.
	Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.

- -