

#3914
Store at -20C**p56Dok-2 Antibody**
Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications: W	Reactivity: H	Sensitivity: Transfected Only	MW (kDa): 56 to 58	Source/Isotype: Rabbit	UniProt ID: #O60496	Entrez-Gene Id: 9046
---------------------------	-------------------------	---	------------------------------	----------------------------------	-------------------------------	--------------------------------

Product Usage Information**Application**

Western Blotting

Dilution

1:1000

Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA and 50% glycerol. Store at -20°C. Do not aliquot the antibody.

Specificity/Sensitivity

p56Dok-2 Antibody detects transfected levels of total p56Dok-2 proteins. The antibody does not cross-react with other p62Dok family members.

Species predicted to react based on 100% sequence homology

Mouse

Source / Purification

Polyclonal antibodies are produced by immunizing animals with a synthetic peptide corresponding to the residues at the carboxy-terminal sequence of human p56Dok-2. The antibodies are purified by protein A and peptide affinity chromatography.

Background

Docking proteins are substrates of tyrosine kinases that function in the recruitment and assembly of specific signal transduction molecules. There are five members in the p62dok family, p62Dok (Dok-1), p56Dok-2 (Dok-2, or DoK-R), Dok-3, Dok-4 and Dok-5 (1-3), characterized by the presence of an amino-terminal PH domain, a central PTB domain and numerous potential sites of tyrosine phosphorylation. Tyrosine phosphorylation of p56Dok-2 occurs upon stimulation of cells with a variety of stimuli, or in cells transformed by oncogenic tyrosine kinases such as v-Src and Bcr-Abl (3-5). Based on the presence of several signaling domains (PH, PTB domain, tyrosine residue and proline-rich regions), it has been proposed that the p62dok family act as docking proteins that link RTKs to signal transduction pathways. p56Dok-2 has been proposed to be a negative regulator of cytokine-induced proliferation in T cells (5). Phosphorylated Tyr351 of p56Dok-2 mediates an association with the SH2 domain of Nck (4).

Background References

1. Master, Z. et al. (2001) *EMBO J.* 20, 5919-5928.
2. Grimm, J. et al. (2001) *J. Cell. Biol.* 154, 345-354.
3. Cristofano, A. D. et al. (1998) *J. Biol. Chem.* 273, 4827-4830.
4. Jones, N. and Dumont, D.J. (1999) *Curr. Biol.* 9, 1057-1060.
5. Nemorin, J.G. and Duplay, P. (2000) *J. Biol. Chem.* 275, 14590-14597.

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

Applications Key**W:** Western Blotting**Cross-Reactivity Key****H:** Human**Trademarks and Patents**

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.**Limited Uses**

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.