

Store at
-20C
#39229**SATB2 (E8R8H) Rabbit mAb**

Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications: W, IP, IHC-P	Reactivity: H	Sensitivity: Endogenous	MW (kDa): 100	Source/Isotype: Rabbit IgG	UniProt ID: #Q9UPW6	Entrez-Gene Id: 23314
--------------------------------------	-------------------------	-----------------------------------	-------------------------	--------------------------------------	-------------------------------	---------------------------------

Product Usage Information**Application**

Western Blotting
Immunoprecipitation
Immunohistochemistry (Paraffin)

Dilution

1:1000
1:50
1:600

Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at -20°C. Do not aliquot the antibody.

Specificity/Sensitivity

SATB2 (E8R8H) Rabbit mAb recognizes endogenous levels of total SATB2 protein. This antibody does not cross-react with SATB1 protein. Non-specific non-nuclear staining was observed in smooth muscle.

Source / Purification

Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to residues surrounding Val476 of human SATB2 protein.

Background

Special AT-rich binding protein 2 (SATB2) is a close homolog to SATB1 that functions as a transcription factor. It binds to nuclear matrix attachment regions (MARS); regulatory DNA sequences important for chromatin structure. SATB2 was initially identified when bound to the MARS of the immunoglobulin µ gene in pre-B cells, enhancing its expression (1). SATB2 plays a role in osteoblast differentiation by repressing the HoxA2 gene and enhancing the activity of Runx2 and ATF4 (2). SATB2 also plays a role in the developing cerebral cortex by changing chromatin structure surrounding the Ctip2 regulatory regions (3). In erythroid cells, SATB2 activates the γ-globin locus by recruiting PCAF and reordering the chromatin structure (4). Downregulation of SATB2 is linked to colorectal cancer and head and neck squamous carcinomas (5,6).

Background References

1. Dobrev, G. et al. (2003) *Genes Dev* 17, 3048-61.
2. Dobrev, G. et al. (2006) *Cell* 125, 971-86.
3. Alcamo, E.A. et al. (2008) *Neuron* 57, 364-77.
4. Zhou, L.Q. et al. (2012) *J Biol Chem* 287, 30641-52.
5. Wang, S. et al. (2009) *J Pathol* 219, 114-22.
6. Liu, T.R. et al. (2012) *PLoS One* 7, e40704.

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

Applications Key

W: Western Blotting **IP:** Immunoprecipitation **IHC-P:** Immunohistochemistry (Paraffin)

Cross-Reactivity Key

H: Human

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

SignalStain is a registered trademark of Cell Signaling Technology, Inc.

XP is a registered trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no

force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.