



Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

Store at -20C
#4078

IκBα (L35A5) Mouse mAb (Amino-terminal Antigen) (Sepharose® Bead Conjugate)

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications: IP	Reactivity: H M R Mk B Pg	Sensitivity: Endogenous	MW (kDa): 39	Source/Isotype: Mouse IgG1	UniProt ID: #P25963	Entrez-Gene Id: 4792
----------------------------	-------------------------------------	-----------------------------------	------------------------	--------------------------------------	-------------------------------	--------------------------------

Product Usage Information

Application

Immunoprecipitation

Dilution

1:20

Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA, 50% glycerol. Store at -20°C. Do not aliquot the antibodies.

Specificity/Sensitivity

IκBα (L35A5) Mouse mAb (Amino-terminal Antigen) detects endogenous levels of total IκBα protein.

Source / Purification

Monoclonal antibody is produced by immunizing animals with a GST-IκBα fusion protein corresponding to the amino-terminus of human IκBα.

Description

This Cell Signaling Technology antibody is immobilized via covalent binding of primary amino groups to N-hydroxysuccinimide (NHS)-activated Sepharose® beads. IκBα (L35A5) Mouse mAb (Amino-terminal Antigen) (Sepharose® Bead Conjugate) is useful for the immunoprecipitation of IκBα protein.

Background

The NF-κB/Rel transcription factors are present in the cytosol in an inactive state complexed with the inhibitory IκB proteins (1-3). Activation occurs via phosphorylation of IκBα at Ser32 and Ser36 followed by proteasome-mediated degradation that results in the release and nuclear translocation of active NF-κB (3-7). IκBα phosphorylation and resulting Rel-dependent transcription are activated by a highly diverse group of extracellular signals including inflammatory cytokines, growth factors, and chemokines. Kinases that phosphorylate IκB at these activating sites have been identified (8).

Background References

1. Baeuerle, P.A. and Baltimore, D. (1988) *Science* 242, 540-6.
2. Beg, A.A. and Baldwin, A.S. (1993) *Genes Dev* 7, 2064-70.
3. Finco, T.S. et al. (1994) *Proc Natl Acad Sci USA* 91, 11884-8.
4. Brown, K. et al. (1995) *Science* 267, 1485-8.
5. Brockman, J.A. et al. (1995) *Mol Cell Biol* 15, 2809-18.
6. Traenckner, E.B. et al. (1995) *EMBO J* 14, 2876-83.
7. Chen, Z.J. et al. (1996) *Cell* 84, 853-62.
8. Karin, M. and Ben-Neriah, Y. (2000) *Annu Rev Immunol* 18, 621-63.

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Applications Key

IP: Immunoprecipitation

Cross-Reactivity Key

H: Human **M:** Mouse **R:** Rat **Mk:** Monkey **B:** Bovine **Pg:** Pig

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for

diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.