## CDCP1 Antibody TECHNOLOGY\* Orders: 877-616-CELL (2355) orders@cellsignal.com



Orders: 877-616-CELL (2355) orders@cellsignal.com Support: 877-678-TECH (8324) Web: info@cellsignal.com cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications: W, IP	<b>Reactivity:</b> H	<b>Sensitivity:</b> Endogenous	<b>MW (kDa):</b> 140, 70	<b>Source/Isotype:</b> Rabbit	UniProt ID: #Q9H5V8	<b>Entrez-Gene Id:</b> 64866
Product Usage Information	2	<b>Application</b> Western Blotting Immunoprecipitation			<b>Dilution</b> 1:1000 1:100	
Storage		Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA and 50% glycerol. Store at – 20°C. Do not aliquot the antibody.				
Specificity/Sensitivity		CDCP1 Antibody detects endogenous levels of total CDCP1 protein. The 140kDa band is full length, glycosylated CDCP1, the 70kDa band is a cleavage product of CDCP1.				
Source / Purification		Polyclonal antibodies are produced by immunizing animals with a synthetic peptide corresponding to amino acids near the carboxy-terminus of human CDCP1. Antibodies are purified by peptide affinity chromatography.				
Background		CUB domain-containing protein 1 (CDCP1, SIMA135) is a putative stem cell marker shown in research studies to be highly expressed in some human cancer cells and in both typical and atypical (cancerous) colons (1). Expression of CDCP1 may be epigenetically regulated, as methylation of promoter CpG sequences results in decreased CDCP1 expression (2). The corresponding <i>CDCP1</i> gene encodes a glycoprotein that acts as a complex, multidomain transmembrane antigen. CDCP1 has three extracellular CUB domains that may be involved in cell adhesion or extracellular matrix interactions (1,3). Src-family kinases may phosphorylate CDCP1 at five tyrosine residues within its cytoplasmic domain to provide a potential binding site for SH2 domain-containing proteins (3). CDCP1 is a putative hematopoietic stem cell marker (4,5).				
Background References		1. Scherl-Mostageer, M 2. Ikeda, J.I. et al. (200 3. Hooper, J.D. et al. (2 4. Conze, T. et al. (200 5. Bühring, H.J. et al. (2	6) <i>J. Pathol.</i> 210, 75- 003) <i>Oncogene</i> 22, 3) <i>Ann. N.Y. Acad. S</i>	84. 1783-94. ci. 996, 222-226.		
Species Reacti	vity	Species reactivity is de	etermined by testing	g in at least one approve	ed application (e.g.,	western blot).
Western Blot Buffer		IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.				
Applications Key		W: Western Blotting IP: Immunoprecipitation				
Cross-Reactivi	ty Key	H: Human				
Trademarks and Patents		Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.				
		All other trademarks a more information.	are the property of t	heir respective owners.	Visit cellsignal.com	/trademarks for
Limited Uses		Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.				
		approved, cleared, or	licensed by the FDA	se Only or a similar labe or other regulatory for duct for any diagnostic o	eign or domestic en	tity, for any

any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products on services that would compete with CST products or services, (c) not alter or remove from the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.