



Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

#41666 Store at +4C

p38 MAPK (D13E1) XP[®] Rabbit mAb (PE Conjugate)

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications: FC-FP	Reactivity: H M R Hm Mk B Pg	Sensitivity: Endogenous	Source/Isotype: Rabbit IgG	UniProt ID: #Q16539, #P53778, #Q15759	Entrez-Gene Id: 1432, 6300, 5600
-------------------------------	--	-----------------------------------	--------------------------------------	--	--

Product Usage Information	Application Flow Cytometry (Fixed/Permeabilized)	Dilution 1:50
Storage	Supplied in PBS (pH 7.2), less than 0.1% sodium azide and 2 mg/ml BSA. Store at 4°C. Do not aliquot the antibody. Protect from light. Do not freeze.	
Specificity/Sensitivity	p38 MAPK (D13E1) XP [®] Rabbit mAb (PE Conjugate) recognizes endogenous levels of total p38 α , β , or γ MAPK protein. This antibody does not recognize p38 δ , SAPK/JNK, or p44/42 MAPK proteins.	
Species predicted to react based on 100% sequence homology	Chicken	
Source / Purification	Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to residues near the carboxy terminus of human p38 protein.	
Description	This Cell Signaling Technology antibody is conjugated to phycoerythrin (PE) and tested in-house for direct flow cytometry analysis in human cells. This antibody is expected to exhibit the same species cross-reactivity as the unconjugated p38 MAPK (D13E1) XP [®] Rabbit mAb #8690.	
Background	p38 MAP kinase (MAPK), also called RK (1) or CSBP (2), is the mammalian orthologue of the yeast HOG kinase that participates in a signaling cascade controlling cellular responses to cytokines and stress (1-4). Four isoforms of p38 MAPK, p38 α , β , γ (also known as Erk6 or SAPK3), and δ (also known as SAPK4) have been identified. Similar to the SAPK/JNK pathway, p38 MAPK is activated by a variety of cellular stresses, including osmotic shock, inflammatory cytokines, lipopolysaccharide (LPS), UV light, and growth factors (1-5). MKK3, MKK6, and SEK activate p38 MAPK by phosphorylation at Thr180 and Tyr182. Activated p38 MAPK has been shown to phosphorylate and activate MAPKAP kinase 2 (3) and to phosphorylate the transcription factors ATF-2 (5), Max (6), and MEF2 (5-8). SB203580 (4-(4-fluorophenyl)-2-(4-methylsulfinylphenyl)-5-(4-pyridyl)-imidazole) is a selective inhibitor of p38 MAPK. This compound inhibits the activation of MAPKAPK-2 by p38 MAPK and subsequent phosphorylation of HSP27 (9). SB203580 inhibits p38 MAPK catalytic activity by binding to the ATP-binding pocket, but does not inhibit phosphorylation of p38 MAPK by upstream kinases (10).	
Background References	<ol style="list-style-type: none"> 1. Rouse, J. et al. (1994) <i>Cell</i> 78, 1027-37. 2. Han, J. et al. (1994) <i>Science</i> 265, 808-11. 3. Lee, J.C. et al. (1994) <i>Nature</i> 372, 739-46. 4. Freshney, N.W. et al. (1994) <i>Cell</i> 78, 1039-49. 5. Raingeaud, J. et al. (1995) <i>J Biol Chem</i> 270, 7420-6. 6. Zervos, A.S. et al. (1995) <i>Proc Natl Acad Sci U S A</i> 92, 10531-4. 7. Zhao, M. et al. (1999) <i>Mol Cell Biol</i> 19, 21-30. 8. Yang, S.H. et al. (1999) <i>Mol Cell Biol</i> 19, 4028-38. 9. Cuenda, A. et al. (1995) <i>FEBS Lett</i> 364, 229-33. 10. Kumar, S. et al. (1999) <i>Biochem Biophys Res Commun</i> 263, 825-31. 	
Species Reactivity	Species reactivity is determined by testing in at least one approved application (e.g., western blot).	
Applications Key	FC-FP: Flow Cytometry (Fixed/Permeabilized)	
Cross-Reactivity Key	H: Human M: Mouse R: Rat Hm: Hamster Mk: Monkey B: Bovine Pg: Pig	
Trademarks and Patents	Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc. XP is a registered trademark of Cell Signaling Technology, Inc.	

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.