



**Orders:** 877-616-CELL (2355)  
orders@cellsignal.com

**Support:** 877-678-TECH (8324)

**Web:** info@cellsignal.com  
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

Store at -20C  
#4194

## Sox1 Antibody

**For Research Use Only. Not for Use in Diagnostic Procedures.**

<b>Applications:</b> W, IF-F	<b>Reactivity:</b> M R	<b>Sensitivity:</b> Endogenous	<b>MW (kDa):</b> 40	<b>Source/Isotype:</b> Rabbit	<b>UniProt ID:</b> #O00570	<b>Entrez-Gene Id:</b> 6656
---------------------------------	---------------------------	-----------------------------------	------------------------	----------------------------------	-------------------------------	--------------------------------

<b>Product Usage Information</b>	<b>Application</b> Western Blotting Immunofluorescence (Frozen)	<b>Dilution</b> 1:1000 1:100 - 1:200
<b>Storage</b>	Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA and 50% glycerol. Store at -20°C. Do not aliquot the antibody.	
<b>Specificity/Sensitivity</b>	Sox1 Antibody detects endogenous levels of total Sox1 protein. It does not cross-react with endogenous levels of other Sox proteins, including Sox2 and Sox3.	
<b>Species predicted to react based on 100% sequence homology</b>	Human	
<b>Source / Purification</b>	Polyclonal antibodies are produced by immunizing animals with a synthetic peptide corresponding to residues at the carboxy terminus of human Sox1. Antibodies are purified by Protein A and peptide affinity chromatography.	
<b>Background</b>	<i>Sox</i> (Sry-related box) genes encode a family of transcription factors that play a myriad of roles in developmental processes (1). Sox1 and Sox2 are the earliest markers that identify neuroectodermal tissue, and these markers, together with Sox9 collectively mark neuronal stem cells (NSC's) that are present in neurogenically active areas of the mature rodent brain (2). Sox1 has been found in the walls of the lateral ventricles and the dentate gyrus, as well as in inactive areas such as the cerebellum where it marks a population of cells in the Purkinje layer known as the Bergmann glia (2). Sox1 is thought to maintain the cell cycle and promote self renewal in NSCs, but also functions in other cell types to promote differentiation (3).	
<b>Background References</b>	<ol style="list-style-type: none"> <li>1. Kiefer, J.C. (2007) <i>Dev Dyn</i> 236, 2356-66.</li> <li>2. Alcock, J. et al. (2009) <i>Neurosci Lett</i> 450, 114-6.</li> <li>3. Kan, L. et al. (2007) <i>Dev Biol</i> 310, 85-98.</li> </ol>	

<b>Species Reactivity</b>	Species reactivity is determined by testing in at least one approved application (e.g., western blot).
<b>Western Blot Buffer</b>	<b>IMPORTANT:</b> For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.
<b>Applications Key</b>	<b>W:</b> Western Blotting <b>IF-F:</b> Immunofluorescence (Frozen)
<b>Cross-Reactivity Key</b>	<b>M:</b> Mouse <b>R:</b> Rat
<b>Trademarks and Patents</b>	Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc. All other trademarks are the property of their respective owners. Visit <a href="http://cellsignal.com/trademarks">cellsignal.com/trademarks</a> for more information.
<b>Limited Uses</b>	Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.  Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any

purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.