## GATA-6 (A549) Antibody



Orders: 877-616-CELL (2355)

orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com

cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

## For Research Use Only. Not for Use in Diagnostic Procedures.

Applications: W	Reactivity: H	<b>Sensitivity:</b> Endogenous	<b>MW (kDa):</b> 55	<b>Source/Isotype:</b> Rabbit	UniProt ID: #Q92908	Entrez-Gene Id: 2627
Product Usage Information		<b>Application</b> Western Blotting			<b>Dilution</b> 1:1000	
Storage		Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 $\mu$ g/ml BSA and 50% glycerol. Store at – 20°C. Do not aliquot the antibody.				
Specificity/Sensitivity		GATA-6 (A549) Antibody detects endogenous levels of total GATA-6 protein.				
Source / Purification		Polyclonal antibodies are produced by immunizing animals with a synthetic peptide corresponding to amino acid sequence surrounding Ala549 of human GATA-6 protein, isoform 1. Antibodies are purified by protein A and peptide affinity chromatography.				
Background		GATA proteins comprise a group of transcription factors that are related by the presence of conserved zinc finger DNA-binding domains, which bind directly to the nucleotide sequence core element GATA (1-3). There are six vertebrate GATA proteins, designated GATA-1 to GATA-6 (3).				
		gut and other organs heart tube and a failu	(5,6). Knock out of re to develop extra	development (4). It is ess GATA-6 is embryonic leth embryonic endoderm (4 ge number of ovarian tu	nal due to defects ir ). Loss of expression	formation of the
Background References		<ol> <li>Ko, L.J. and Engel, J.D. (1993) <i>Mol Cell Biol</i> 13, 4011-22.</li> <li>Merika, M. and Orkin, S.H. (1993) <i>Mol Cell Biol</i> 13, 3999-4010.</li> <li>Lowry, J.A. and Atchley, W.R. (2000) <i>J Mol Evol</i> 50, 103-15.</li> <li>Cai, K.Q. et al. (2008) <i>Dev Dyn</i> 237, 2820-9.</li> <li>Charron, F. and Nemer, M. (1999) <i>Semin Cell Dev Biol</i> 10, 85-91.</li> <li>Haveri, H. et al. (2008) <i>BMC Gastroenterol</i> 8, 9.</li> <li>Caslini, C. et al. (2006) <i>Oncogene</i> 25, 5446-61.</li> </ol>				
Species Reactiv	rity	Species reactivity is de	etermined by testin	g in at least one approve	ed application (e.g.,	western blot).
Western Blot Buffer		IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X				

Western Blot Buffer IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X

TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

Applications Key W: Western Blotting

Cross-Reactivity Key H: Human

**Trademarks and Patents** Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for

more information.

**Limited Uses**Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Broducts provided by CST, its affiliates or its distributors. Any Customer's

the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no

force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for

Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.