

#42847  
Store at +4C**CD11c (3.9) Mouse mAb (redFluor™ 710 Conjugate)**

**Orders:** 877-616-CELL (2355)  
orders@cellsignal.com

**Support:** 877-678-TECH (8324)

**Web:** info@cellsignal.com  
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

**For Research Use Only. Not for Use in Diagnostic Procedures.**

<b>Applications:</b> FC-FP, FC-L	<b>Reactivity:</b> H	<b>Sensitivity:</b> Endogenous	<b>Source/Isotype:</b> Mouse IgG1	<b>UniProt ID:</b> #P20702	<b>Entrez-Gene Id:</b> 3687
-------------------------------------	-------------------------	-----------------------------------	--------------------------------------	-------------------------------	--------------------------------

<b>Product Usage Information</b>	<b>Application</b> Flow Cytometry (Fixed/Permeabilized) Flow Cytometry (Live)	<b>Dilution</b> 1:20 1:20
<b>Storage</b>	Supplied in 10 mM NaH <sub>2</sub> PO <sub>4</sub> , 150 mM NaCl, 0.09% NaN <sub>3</sub> , 0.1% gelatin, pH 7.2. This product is stable for 12 months when stored at 4°C. Do not aliquot the antibody. Protect from light. Do not freeze.	
<b>Specificity/Sensitivity</b>	CD11c (3.9) Mouse mAb (redFluor™ 710 Conjugate) recognizes endogenous levels of total CD11c protein. This antibody detects an epitope within the extracellular domain.	
<b>Source / Purification</b>	This monoclonal antibody was purified from tissue culture supernatant via affinity chromatography. The purified antibody was conjugated under optimal conditions, with unreacted dye removed from the preparation.	
<b>Description</b>	This Cell Signaling Technology antibody is conjugated to redFluor™ 710 and tested in-house for direct flow cytometry analysis in human cells.	
<b>Background</b>	CD11c (integrin $\alpha$ X, ITGAX) is a transmembrane glycoprotein that forms an $\alpha\beta$ heterodimer with CD18 (integrin $\beta$ 2), which interacts with a variety of extracellular matrix molecules and cell surface proteins (1). CD11c is primarily used as a dendritic cell marker. Dendritic cells can be classified into two major types: CD11c+ conventional dendritic cells that specialize in antigen presentation, and CD11c- plasmacytoid dendritic cells that specialize in type I interferon production (2, 3). CD11c expression has also been observed on activated NK cells, subsets of B cells, monocytes, granulocytes, and some B cell malignancies including hairy cell leukemia (4-7). The 3.9 antibody is widely used as a marker for CD11c expression on the above mentioned cell types.	
<b>Background References</b>	<ol style="list-style-type: none"> <li>1. Uotila, L.M. et al. (2013) <i>J Biol Chem</i> 288, 33494-9.</li> <li>2. Kohrgruber, N. et al. (1999) <i>J Immunol</i> 163, 3250-9.</li> <li>3. Siegal, F.P. et al. (1999) <i>Science</i> 284, 1835-7.</li> <li>4. Racine, R. et al. (2008) <i>J Immunol</i> 181, 1375-85.</li> <li>5. Werfel, T. et al. (1991) <i>J Immunol</i> 147, 2423-7.</li> <li>6. Cabañas, C. et al. (1988) <i>Hybridoma</i> 7, 167-76.</li> <li>7. Kristensen, J.S. et al. (1987) <i>Blood</i> 70, 1063-8.</li> </ol>	
<b>Species Reactivity</b>	Species reactivity is determined by testing in at least one approved application (e.g., western blot).	
<b>Applications Key</b>	<b>FC-FP:</b> Flow Cytometry (Fixed/Permeabilized) <b>FC-L:</b> Flow Cytometry (Live)	
<b>Cross-Reactivity Key</b>	<b>H:</b> Human	
<b>Trademarks and Patents</b>	Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.  All other trademarks are the property of their respective owners. Visit <a href="http://cellsignal.com/trademarks">cellsignal.com/trademarks</a> for more information.	
<b>Limited Uses</b>	<p>Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.</p> <p>Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any</p>	

purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.