

UGT Antibody



Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications:	Reactivity:	Sensitivity:	MW (kDa):	Source/Isotype:	UniProt ID:	Entrez-Gene Id:
W	H M R	Endogenous	55-60	Rabbit	#Q9HAW8, #O60656, #P35503, #P22309, #P35504, #P22310, #Q9HAW7, #Q9HAW9, #P19224	54575, 54600, 54659, 54658, 54579, 54657, 54577, 54576, 54578

Product Usage Information	Application	Dilution
Storage	Western Blotting	1:1000
Specificity/Sensitivity	Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA and 50% glycerol. Store at -20°C. Do not aliquot the antibody.	
Source / Purification	UGT Antibody detects endogenous levels of proteins from the UGT1A subfamily.	
Background	Polyclonal antibodies are produced by immunizing animals with a synthetic peptide corresponding to residues near the carboxy terminus of the UGT1A subfamily. Antibodies were purified by protein A and peptide affinity chromatography.	
Background References	Glucuronidation is a major pathway that enhances the elimination of lipophilic xenobiotics and endobiotics to more more water soluble compounds for excretion (1,2). The UDP-glucuronosyltransferase (UGT) superfamily catalyzes the glucuronidation of the glycosyl group of a nucleotide sugar to a variety of endogenous and exogenous compounds. Over 100 UGT mammalian gene products have been described and have been divided into subfamilies based on sequence identities (3). The UGT1 subfamily consists of a number of gene products resulting from alternative splicing. These UGT products can differ in tissue expression and substrate specificity. Also, marked differences in the individual expression of UGT isoforms can account for differences in drug metabolism.	
	<ol style="list-style-type: none"> 1. Wells, P.G. et al. (2004) <i>Drug Metab. Dispos.</i> 32, 281-90. 2. King, C.D. et al. (2000) <i>Curr. Drug Metab.</i> 1, 143-61. 3. Mackenzie, P.I. et al. (2005) <i>Pharmacogenet. Genomics</i> 15, 677-85. 	

Species Reactivity	Species reactivity is determined by testing in at least one approved application (e.g., western blot).
Western Blot Buffer	IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween@ 20 at 4°C with gentle shaking, overnight.
Applications Key	W: Western Blotting
Cross-Reactivity Key	H: Human M: Mouse R: Rat
Trademarks and Patents	Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc. All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.
Limited Uses	Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.