



**Orders:** 877-616-CELL (2355)  
orders@cellsignal.com

**Support:** 877-678-TECH (8324)

**Web:** info@cellsignal.com  
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

Store at -20C  
#4392

## PIAS4 (D2F12) Rabbit mAb

For Research Use Only. Not for Use in Diagnostic Procedures.

<b>Applications:</b> W	<b>Reactivity:</b> H R Mk	<b>Sensitivity:</b> Endogenous	<b>MW (kDa):</b> 75	<b>Source/Isotype:</b> Rabbit IgG	<b>UniProt ID:</b> #Q8N2W9	<b>Entrez-Gene Id:</b> 51588
---------------------------	------------------------------	-----------------------------------	------------------------	--------------------------------------	-------------------------------	---------------------------------

### Product Usage Information

#### Application

Western Blotting

#### Dilution

1:1000

### Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at -20°C. Do not aliquot the antibody.

### Specificity/Sensitivity

PIAS4 (D2F12) Rabbit mAb detects endogenous levels of total PIAS4 protein.

### Species predicted to react based on 100% sequence homology

Mouse

### Source / Purification

Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to residues surrounding Lys59 of human PIAS4 protein.

### Background

The protein inhibitor of activated Stat (PIAS) proteins, which include PIAS1, PIAS3, PIASx, and PIASy, were originally characterized based on their interaction with the Stat family of transcription factors (1,2). PIAS1, PIAS3, and PIASx interact with and repress Stat1, Stat3, and Stat4, respectively (1-3). Deletion of PIAS1 leads to inhibition of interferon-inducible genes and increased protection against infection (4). The PIAS family contains a conserved RING domain that has been linked to a function as a small ubiquitin-related modifier (SUMO) ligase, coupling the SUMO conjugating enzyme Ubc9 with its substrate proteins (5,6). Numerous studies have now shown that PIAS family members can regulate the activity of transcription factors through distinct mechanisms, including NF-κB (7,8), c-Jun, p53 (5,9), Oct-4 (10), and Smads (11,12). The activity of PIAS1 is regulated by both phosphorylation and arginine methylation. Inflammatory stimuli can induce IKK-mediated phosphorylation of PIAS1 at Ser90, which is required for its activity (13). In addition, PRMT1 induces arginine methylation of PIAS1 at Arg303 following interferon treatment and is associated with its repressive activity on Stat1 (14). PIAS4, also known as PIASy, is a specific SUMO-E3 ligase for Ets-1 and represses Ets-1 dependent transcription (15). PIAS4 also alters the nuclear localization, reduces the transcriptional activity of C/EBPδ, and enhances cell proliferation and migration (16).

### Background References

- Liu, B. et al. (1998) *Proc Natl Acad Sci USA* 95, 10626-31.
- Chung, C.D. et al. (1997) *Science* 278, 1803-5.
- Arora, T. et al. (2003) *J Biol Chem* 278, 21327-30.
- Liu, B. et al. (2004) *Nat Immunol* 5, 891-8.
- Schmidt, D. and Müller, S. (2002) *Proc Natl Acad Sci USA* 99, 2872-7.
- Kotaja, N. et al. (2002) *Mol Cell Biol* 22, 5222-34.
- Liu, B. et al. (2005) *Mol Cell Biol* 25, 1113-23.
- Tahk, S. et al. (2007) *Proc Natl Acad Sci USA* 104, 11643-8.
- Bischof, O. et al. (2006) *Mol Cell* 22, 783-94.
- Tolkunova, E. et al. (2007) *J Mol Biol* 374, 1200-12.
- Long, J. et al. (2004) *Proc Natl Acad Sci USA* 101, 99-104.
- Murdoch, R.N. and Edwards, T. (1992) *Biochem Int* 28, 1029-37.
- Liu, B. et al. (2007) *Cell* 129, 903-14.
- Weber, S. et al. (2009) *Genes Dev* 23, 118-32.
- Nishida, T. et al. (2007) *Biochem J* 405, 481-8.
- Zhou, S. et al. (2008) *J Biol Chem* 283, 20137-48.

### Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

### Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

**Applications Key****W:** Western Blotting**Cross-Reactivity Key****H:** Human **R:** Rat **Mk:** Monkey**Trademarks and Patents**

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit [cellsignal.com/trademarks](http://cellsignal.com/trademarks) for more information.**Limited Uses**

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.