CaMKII (pan) (D11A10) Rabbit mAb





Orders:	877-616-CELL (2355) orders@cellsignal.com			
Support:	877-678-TECH (8324)			
Web:	info@cellsignal.com cellsignal.com			

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications: W	Reactivity: H M R	Sensitivity: Endogenous	MW (kDa): 60, 50	Source/Isotype: Rabbit IgG	UniProt ID: #Q13554, #Q13557, #Q13555, #Q9UQM7	Entrez-Gene Id: 816, 817, 818, 815		
Product Usage Information		Application Western Blotting			Dilution 1:1000			
Storage		Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 μg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at –20°C. Do not aliquot the antibody.						
Specificity/Sen	sitivity	CaMKII (pan) (D11A10) Rabbit mAb detects endogenous levels of total CaMKII protein. The peptide sequence used as the antigen is 100% conserved between CaMKII-alpha, gamma and delta, and 88% conserved in CaMKII-beta.						
Source / Purifi	urce / Purification Monoclonal antibody is produced by immunizing animals with a synthetic peptide surroundin of human CaMKII-alpha.							
Background	undCaMKII is an important member of the calcium/calmodulin-activated protein kinase family, function in neural synaptic stimulation and T cell receptor signaling (1,2). CaMKII has catalytic and regulator domains. Ca ²⁺ /calmodulin binding to the CaMKII regulatory domain relieves autoinhibition and activates the kinase (3). The activated CaMKII further autophosphorylates at Thr286 to render the kinase constitutively active (3). The threonine phosphorylation state of CaMKII can be regulated through PP1/PKA. PP1 (protein phosphatase 1) dephosphorylates phospho-CaMKII at Thr286. PKA (protein kinase A) prevents phospho-CaMKII (Thr286) dephosphorylation through an inhibitory effe on PP1 (4).							
Background Re	eferences	1. Hughes, K. et al. (20 2. Barria, A. et al. (199 3. Barkai, U. et al. (200 4. Makhinson, M. et al	7) <i>Science</i> 276, 2042 0) <i>Mol Endocrinol</i> 1	2-5. 4, 554-63.				
Species Reactiv	vity	Species reactivity is de	etermined by testing	g in at least one approve	d application (e.g.,	, western blot).		
Western Blot BufferIMPORTANT: For western blots, incubate membrane with diluted p TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.					primary antibody i	n 5% w/v BSA, 1X		
Applications K	ey	W: Western Blotting						
Cross-Reactivit	ту Кеу	H: Human M: Mouse R: Rat						
Trademarks and Patents		Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.						
		All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.						
Limited Uses		Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.						
		Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in						

any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products on services that would compete with CST products or services, (c) not alter or remove from the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.