



Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

Store at -20C
#4806

Phospho-NF- κ B p105 (Ser932) (18E6) Rabbit mAb

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications:	Reactivity:	Sensitivity:	MW (kDa):	Source/Isotype:	UniProt ID:	Entrez-Gene Id:
W, IP	H M R Mk	Endogenous	120	Rabbit IgG	#P19838	4790

Product Usage Information

Application

Western Blotting
Immunoprecipitation

Dilution

1:1000
1:100

Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 μ g/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at -20°C . Do not aliquot the antibody.

Specificity/Sensitivity

Phospho-NF- κ B p105 (Ser932) (18E6) Rabbit mAb detects endogenous levels of p105NF- κ B only when phosphorylated at serine 932.

Source / Purification

Monoclonal antibody is produced by immunizing animals with a synthetic phosphopeptide corresponding to amino acids around Ser932 of NF- κ B p105.

Background

Transcription factors of the nuclear factor κ B (NF- κ B)/Rel family play a pivotal role in inflammatory and immune responses (1,2). There are five family members in mammals: RelA, c-Rel, RelB, NF- κ B1 (p105/p50), and NF- κ B2 (p100/p52). Both p105 and p100 are proteolytically processed by the proteasome to produce p50 and p52, respectively. Rel proteins bind p50 and p52 to form dimeric complexes that bind DNA and regulate transcription. In unstimulated cells, NF- κ B is sequestered in the cytoplasm by I κ B inhibitory proteins (3-5). NF- κ B-activating agents can induce the phosphorylation of I κ B proteins, targeting them for rapid degradation through the ubiquitin-proteasome pathway and releasing NF- κ B to enter the nucleus where it regulates gene expression (6-8). NIK and IKK α (IKK1) regulate the phosphorylation and processing of NF- κ B2 (p100) to produce p52, which translocates to the nucleus (9-11).

Following IKK-mediated phosphorylation of p105 NF- κ B at multiple sites (Ser921, 923, 927, and 932) on its carboxy-terminus, SCF β -TrCP mediated processing produces the 50kDa active form p50 (12,13).

Background References

- Baeuerle, P.A. and Henkel, T. (1994) *Annu Rev Immunol* 12, 141-79.
- Baeuerle, P.A. and Baltimore, D. (1996) *Cell* 87, 13-20.
- Haskill, S. et al. (1991) *Cell* 65, 1281-9.
- Thompson, J.E. et al. (1995) *Cell* 80, 573-82.
- Whiteside, S.T. et al. (1997) *EMBO J* 16, 1413-26.
- Traenckner, E.B. et al. (1995) *EMBO J* 14, 2876-83.
- Scherer, D.C. et al. (1995) *Proc Natl Acad Sci USA* 92, 11259-63.
- Chen, Z.J. et al. (1996) *Cell* 84, 853-62.
- Senftleben, U. et al. (2001) *Science* 293, 1495-9.
- Coope, H.J. et al. (2002) *EMBO J* 21, 5375-85.
- Xiao, G. et al. (2001) *Mol Cell* 7, 401-9.
- Heissmeyer, V. et al. (2001) *Mol Cell Biol* 21, 1024-35.
- Orian, A. et al. (2000) *EMBO J* 19, 2580-91.

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween@ 20 at 4 $^{\circ}$ C with gentle shaking, overnight.

Applications Key

W: Western Blotting **IP:** Immunoprecipitation

Cross-Reactivity Key

H: Human **M:** Mouse **R:** Rat **Mk:** Monkey

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.