

#4820 Store at -20C

CREB (D76D11) Rabbit mAb



Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications:	Reactivity:	Sensitivity:	MW (kDa):	Source/Isotype:	UniProt ID:	Entrez-Gene Id:
W, W-S, IP, IF-F, IF-IC, FC-FP, ChIP, C&T	H M R Hm Mk Dm	Endogenous	43	Rabbit IgG	#P16220	1385

Product Usage Information

For optimal ChIP results, use 5 µl of antibody and 10 µg of chromatin (approximately 4 x 10⁶ cells) per IP. This antibody has been validated using SimpleChIP[®] Enzymatic Chromatin IP Kits.

Application	Dilution
Western Blotting	1:1000
Simple Western™	1:50 - 1:250
Immunoprecipitation	1:50
Immunofluorescence (Frozen)	1:400 - 1:1600
Immunofluorescence (Immunocytochemistry)	1:400 - 1:1600
Flow Cytometry (Fixed/Permeabilized)	1:200 - 1:800
Chromatin IP	1:100
CUT&Tag	1:50

Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at -20°C. Do not aliquot the antibody.

Specificity/Sensitivity

CREB (D76D11) Rabbit mAb detects endogenous levels of total CREB-1 protein. The antibody does not cross-react with other ATF/CREB family members.

Source / Purification

Monoclonal antibody is produced by immunizing animals with a full length GST-CREB fusion protein. The epitope has been mapped to residues surrounding Gly61.

Background

CREB is a bZIP transcription factor that activates target genes through cAMP response elements. CREB is able to mediate signals from numerous physiological stimuli, resulting in regulation of a broad array of cellular responses. While CREB is expressed in numerous tissues, it plays a large regulatory role in the nervous system. CREB is believed to play a key role in promoting neuronal survival, precursor proliferation, neurite outgrowth, and neuronal differentiation in certain neuronal populations (1-3). Additionally, CREB signaling is involved in learning and memory in several organisms (4-6). CREB is able to selectively activate numerous downstream genes through interactions with different dimerization partners. CREB is activated by phosphorylation at Ser133 by various signaling pathways, including Erk, Ca²⁺, and stress signaling. Some of the kinases involved in phosphorylating CREB at Ser133 are p90RSK, MSK, CaMKIV, and MAPKAPK-2 (7-9).

Background References

1. Lonze, B.E. et al. (2002) *Neuron* 34, 371-85.
2. Lee, M.M. et al. (1999) *J Neurosci Res* 55, 702-12.
3. Redmond, L. et al. (2002) *Neuron* 34, 999-1010.
4. Dash, P.K. et al. (1990) *Nature* 345, 718-21.
5. Yin, J.C. et al. (1994) *Cell* 79, 49-58.
6. Guzowski, J.F. and McGaugh, J.L. (1997) *Proc Natl Acad Sci USA* 94, 2693-8.
7. Xing, J. et al. (1998) *Mol Cell Biol* 18, 1946-55.
8. Ribar, T.J. et al. (2000) *J Neurosci* 20, RC107.
9. Tan, Y. et al. (1996) *EMBO J* 15, 4629-42.

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

Applications Key

W: Western Blotting **W-S:** Simple Western™ **IP:** Immunoprecipitation **IF-F:** Immunofluorescence (Frozen) **IF-IC:** Immunofluorescence (Immunocytochemistry) **FC-FP:** Flow Cytometry

(Fixed/Permeabilized) **ChIP**: Chromatin IP **C&T**: CUT&Tag

Cross-Reactivity Key

H: Human **M**: Mouse **R**: Rat **Hm**: Hamster **Mk**: Monkey **Dm**: D. melanogaster

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.