

## Hic-5 Antibody



**Orders:** 877-616-CELL (2355)  
orders@cellsignal.com

**Support:** 877-678-TECH (8324)

**Web:** info@cellsignal.com  
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications:	Reactivity:	Sensitivity:	MW (kDa):	Source/Isotype:	UniProt ID:	Entrez-Gene Id:
W, IP	H Mk B	Endogenous	50	Rabbit	#O43294	7041

### Product Usage Information

#### Application

Western Blotting  
Immunoprecipitation

#### Dilution

1:1000  
1:100

### Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA and 50% glycerol. Store at -20°C. Do not aliquot the antibody.

### Specificity/Sensitivity

Hic-5 Antibody detects endogenous levels of total Hic-5/ARA55 protein.

### Source / Purification

Polyclonal antibodies are produced by immunizing animals with a synthetic peptide corresponding to residues surrounding Ala104 of human Hic-5. Antibodies are purified by protein A and peptide affinity chromatography.

### Background

Hic-5 is a LIM domain family member originally identified as a TGFβ1 and hydrogen peroxide inducible gene, and is nearly identical to the androgen receptor co-activator ARA55 (1-3). Hic-5 is structurally related to paxillin, and both proteins are localized to focal adhesions and thought to serve as adaptor molecules, linking signals from the extracellular matrix to cytoskeletal regulation and intracellular signaling (4,5). Like paxillin, Hic-5 contains four LD motifs and four LIM domains. Expression of Hic-5 can affect cell growth and differentiation (6-8). Increased expression of Hic-5 is observed during cellular senescence in fibroblasts, and ectopic expression in immortalized fibroblasts suppressed cell growth (8). Unlike paxillin, Hic-5 may translocate to the nucleus in response to oxidants like hydrogen peroxide (9). It has been proposed that Hic-5 serves to shuttle redox signaling from focal adhesions to the nucleus where it acts as a transcriptional co-activator for some transcription factors including, Sp1 and PPARγ (7,9,10). Phosphorylation of Hic-5 at Tyr60 by CAKβ and Fyn may activate Hic-5 signaling by allowing binding to downstream SH2 domain containing proteins (11).

### Background References

- Ohba, M. et al. (1994) *J Cell Biol* 126, 1079-88.
- Shibanuma, M. et al. (1994) *J Biol Chem* 269, 26767-74.
- Fujimoto, N. et al. (1999) *J Biol Chem* 274, 8316-21.
- Matsuya, M. et al. (1998) *J Biol Chem* 273, 1003-14.
- Nishiya, N. et al. (2001) *Mol Cell Biol* 21, 5332-45.
- Hu, Y. et al. (1999) *Proc Natl Acad Sci U S A* 96, 10218-23.
- Drori, S. et al. (2005) *Genes Dev* 19, 362-75.
- Shibanuma, M. et al. (1997) *Mol Cell Biol* 17, 1224-35.
- Shibanuma, M. et al. (2003) *Mol Biol Cell* 14, 1158-71.
- Shibanuma, M. et al. (2004) *J Cell Biochem* 91, 633-45.
- Ishino, M. et al. (2000) *FEBS Lett* 474, 179-83.

### Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

### Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

### Applications Key

**W:** Western Blotting **IP:** Immunoprecipitation

### Cross-Reactivity Key

**H:** Human **Mk:** Monkey **B:** Bovine

### Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit [cellsignal.com/trademarks](http://cellsignal.com/trademarks) for more information.

## Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.