

Phospho-MEK1 (Thr292) Antibody

Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications:	Reactivity:	Sensitivity:	MW (kDa):	Source/Isotype:	UniProt ID:	Entrez-Gene Id:
W, IP	H M R Mk	Endogenous	45	Rabbit	#Q02750	5604

Product Usage Information**Application**

Western Blotting
Immunoprecipitation

Dilution

1:1000
1:200

Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA and 50% glycerol. Store at -20°C. Do not aliquot the antibody.

Specificity/Sensitivity

Phospho-MEK1 (Thr292) Antibody recognizes endogenous levels of MEK1 protein only when phosphorylated at Thr292. This antibody does not cross-react with MEK2 protein. This antibody cross-reacts with a 170 kDa protein of unknown identity in some cell lysates.

Species predicted to react based on 100% sequence homology

Dog, Pig

Source / Purification

Polyclonal antibodies are produced by immunizing animals with a synthetic phosphopeptide corresponding to residues surrounding Thr292 of human MEK1 protein. Antibodies are purified by protein A and peptide affinity chromatography.

Background

MEK1 and MEK2, also called MAPK or Erk kinases, are dual-specificity protein kinases that function in a mitogen activated protein kinase cascade controlling cell growth and differentiation (1-3). Activation of MEK1 and MEK2 occurs through phosphorylation of two serine residues at positions 217 and 221, located in the activation loop of subdomain VIII, by Raf-like molecules. MEK1/2 is activated by a wide variety of growth factors and cytokines and also by membrane depolarization and calcium influx (1-4). Constitutively active forms of MEK1/2 are sufficient for the transformation of NIH/3T3 cells or the differentiation of PC-12 cells (4). MEK activates p44 and p42 MAP kinase by phosphorylating both threonine and tyrosine residues at sites located within the activation loop of kinase subdomain VIII. In response to integrin signaling, p21-activated kinase -1 (PAK-1) phosphorylates MEK1 at Ser298, which enhances MEK1-ERK2 complex formation and MEK1 activation by Raf-1. These events positively regulate the Raf-1-MEK-ERK signaling cascade (5-9). Research studies have identified a negative regulatory loop in the Raf-1-MEK-ERK signaling cascade, in which ERK2-dependent phosphorylation of MEK1 at Thr292 negatively regulates MEK1 activation following cell adhesion. Specifically, ERK2-dependent phosphorylation of MEK1 also attenuates its PAK-1-mediated phosphorylation, contributing to a reduction in Raf-dependent phosphorylation of residues within the MEK1 activation loop (7,10).

Background References

1. Crews, C.M. et al. (1992) *Science* 258, 478-480.
2. Alessi, D.R. et al. (1994) *EMBO J.* 13, 1610-19.
3. Rosen, L.B. et al. (1994) *Neuron* 12, 1207-21.
4. Cowley, S. et al. (1994) *Cell* 77, 841-52.
5. Frost, J.A. et al. (1997) *EMBO J* 16, 6426-38.
6. Rossomando, A.J. et al. (1994) *Mol Cell Biol* 14, 1594-602.
7. Eblen, S.T. et al. (2004) *Mol Cell Biol* 24, 2308-17.
8. Eblen, S.T. et al. (2002) *Mol Cell Biol* 22, 6023-33.
9. Coles, L.C. and Shaw, P.E. (2002) *Oncogene* 21, 2236-44.
10. Xu, B.e. et al. (1999) *J Biol Chem* 274, 34029-35.

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v nonfat dry milk, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

Applications Key

W: Western Blotting **IP:** Immunoprecipitation

Cross-Reactivity Key

H: Human **M:** Mouse **R:** Rat **Mk:** Monkey

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

XP is a registered trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.

Orders: 877-616-CELL (2355) • orders@cellsignal.com • Support: 877-678-TECH (8324) • info@cellsignal.com • Web: cellsignal.com
For Research Use Only. Not for Use in Diagnostic Procedures.