

**MLL3 (D1S1V) Rabbit mAb**

**Orders:** 877-616-CELL (2355)  
orders@cellsignal.com

**Support:** 877-678-TECH (8324)

**Web:** info@cellsignal.com  
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

**For Research Use Only. Not for Use in Diagnostic Procedures.**

<b>Applications:</b> W	<b>Reactivity:</b> H	<b>Sensitivity:</b> Endogenous	<b>MW (kDa):</b> 540	<b>Source/Isotype:</b> Rabbit IgG	<b>UniProt ID:</b> #Q8NEZ4	<b>Entrez-Gene Id:</b> 58508
---------------------------	-------------------------	-----------------------------------	-------------------------	--------------------------------------	-------------------------------	---------------------------------

**Product Usage Information****Application**

Western Blotting

**Dilution**

1:1000

**Storage**

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at -20°C. Do not aliquot the antibody.

**Specificity/Sensitivity**

MLL3 (D1S1V) Rabbit mAb recognizes endogenous levels of total MLL3 protein.

**Source / Purification**

Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to residues surrounding Ala2110 of human MLL3 protein.

**Background**

The Set1 histone methyltransferase protein was first identified in yeast as part of the Set1/COMPASS histone methyltransferase complex, which methylates histone H3 at Lys4 and functions as a transcriptional co-activator (1). While yeast contain only one known Set1 protein, mammals contain six Set1-related proteins: SET1A, SET1B, MLL1, MLL2, MLL3, and MLL4, all of which assemble into COMPASS-like complexes and methylate histone H3 at Lys4 (2,3). These Set1-related proteins are each found in distinct protein complexes, all of which share the common subunits WDR5, RBBP5, ASH2L, CXXC1 and DPY30, which are required for proper complex assembly and modulation of histone methyltransferase activity (2-6). MLL1 and MLL2 complexes contain the additional protein subunit, menin (6).

MLL3, also known as histone-lysine N-methyltransferase 2C (KMT2C), is a large 540 kDa protein that functions as part of the MLL3/COMPASS-like complex to activate gene expression by mediating mono-methylation of histone H3 lysine 4 at gene enhancers (7). Enhancer-specific H3 lysine 4 mono-methylation (H3K4me1) correlates with increased levels of chromatin interactions between gene enhancers and promoters, while loss of this modification results in a reduction of enhancer-promoter interactions (8). Furthermore, H3K4me1 facilitates recruitment of the Cohesin complex, which may function to promote the interactions between gene enhancers and promoters (8). MLL3 is found to be mutated or have altered expression in a number of different cancers (9).

**Background References**

1. Miller, T. et al. (2001) *Proc Natl Acad Sci U S A* 98, 12902-7.
2. Shilatifard, A. (2008) *Curr Opin Cell Biol* 20, 341-8.
3. Tenney, K. and Shilatifard, A. (2005) *J Cell Biochem* 95, 429-36.
4. Lee, J.H. and Skalnik, D.G. (2005) *J Biol Chem* 280, 41725-31.
5. Lee, J.H. et al. (2007) *J Biol Chem* 282, 13419-28.
6. Hughes, C.M. et al. (2004) *Mol Cell* 13, 587-97.
7. Hu, D. et al. (2013) *Mol Cell Biol* 33, 4745-54.
8. Yan, J. et al. (2018) *Cell Res*, .
9. Sze, C.C. and Shilatifard, A. (2016) *Cold Spring Harb Perspect Med* 6, .

**Species Reactivity**

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

**Western Blot Buffer**

**IMPORTANT:** For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

**Applications Key**

**W:** Western Blotting

**Cross-Reactivity Key**

**H:** Human

**Trademarks and Patents**

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.  
XP is a registered trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit [cellsignal.com/trademarks](http://cellsignal.com/trademarks) for more information.

## Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.