## 45419

## ZFX (L28B6) Mouse mAb



Orders: 877-616-CELL (2355)

orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com

cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

## For Research Use Only. Not for Use in Diagnostic Procedures.

Applications: W	Reactivity: H	<b>Sensitivity:</b> Endogenous	<b>MW (kDa):</b> 135	<b>Source/Isotype:</b> Mouse IgG2b	UniProt ID: #P17010	Entrez-Gene Id: 7543
Product Usage Information		<b>Application</b> Western Blotting	<b>Dilution</b> 1:1000			
Storage		Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 μg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at –20°C. Do not aliquot the antibody.				
Specificity/Sensitivity		ZFX (L28B6) Mouse mAb detects endogenous levels of total ZFX protein. This antibody may cross-react with ZFY protein.				
Source / Purification		Monoclonal antibody is produced by immunizing animals with a recombinant fragment of human ZFX.				
Background		The ZFX (X-linked zinc finger protein) gene is expressed from the inactive X chromosome and is structurally similar to its homologue on the Y chromosome, ZFY (1). Transcripts of ZFX and ZFY encode proteins composed of a highly acidic amino-terminal domain and a carboxy-terminal zinc-finger motif that is commonly associated with nucleic acid-binding proteins (2). Both ZFY and ZFX are probable transcriptional activators and may function in sex determination (3). Multiple alternatively spliced transcript variants of ZFX, encoding different isoforms, have been identified and may be functionally distinct (2). Conditional gene targeting studies in mouse have suggested ZFX is also required for self-renewal of embryonic and hematopoietic stem cells (4). ZFX has also been suggested to play a role in proliferation and expansion of B cells, and could contribute to lymphocyte homeostasis (5).				
Background References		1. Schneider-Gädicke, A. et al. (1989) <i>Cell</i> 57, 1247-58. 2. Schneider-Gädicke, A. et al. (1989) <i>Nature</i> 342, 708-11. 3. Palmer, M.S. et al. (1990) <i>Proc Natl Acad Sci U S A</i> 87, 1681-5. 4. Galan-Caridad, J.M. et al. (2007) <i>Cell</i> 129, 345-57. 5. Arenzana, T.L. et al. (2009) <i>Blood</i> 113, 5857-67.				
Species Reactivity		Species reactivity is determined by testing in at least one approved application (e.g., western blot).				
Western Blot Buffer		IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.				
Applications Key		W: Western Blotting				
Cross-Reactivity Key		H: Human				
Trademarks and Patents		Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.				
		All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.				
Limited Uses		the following terms a terms and conditions	wise expressly agreed in a writing signed by a legally authorized representative of CST, erms apply to Products provided by CST, its affiliates or its distributors. Any Customer's ditions that are in addition to, or different from, those contained herein, unless expected in writing by a legally authorized representative of CST, are rejected and are of no			

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a

component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.