

29*V*57

Phospho-Tyrosine Mouse mAb (P-Tyr-100) (HRP Conjugate)



Orders: 877-616-CELL (2355)

orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com

cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications: W, E-P	Reactivity: All	Sensitivity: Endogenous	Source/Isotype: Mouse IgG1	
Product Usage Information		Application Western Blotting Peptide ELISA (DELFIA)	Dilution 1:2000 1:1000	
Storage			3 mM KCI, 10 mM sodium phosphate (pH 7.4) dibasic, 2 mM potassium mg/mL BSA, and 50% glycerol. Store at –20°C. <i>Do not aliquot the antibody.</i>	
Specificity/Sensitivity		Phospho-Tyrosine Mouse mAb (P-Tyr-100) (HRP Conjugate) is a high affinity antibody. ELISAs against a wide variety of phosphopeptides indicate that P-Tyr-100 binds phospho-Tyr in a manner largely independent of the surrounding amino acid sequence. 2D gel western blot analysis of pervanadate-treated cell extracts also shows that P-Tyr-100 interacts with a broad range of tyrosine-phosphorylated proteins. P-Tyr-100 does not cross-react with peptides containing phospho-Ser or phospho-Thr. (U.S. Patent No's.: 6,441,140; 6,982,318; 7,259,022; 7,344,714; U.S.S.N. 11,484,485; and all foreign equivalents.)		
Source / Purification		Monoclonal antibody is produced by immunizing animals with phospho-tyrosine containing peptides.		
Description		This Cell Signaling Technology [®] antibody is conjugated to the carbohydrate groups of horseradish peroxidase (HRP) via its amine groups. The HRP conjugated antibody is expected to exhibit the same species cross-reactivity as the unconjugated Phospho-Tyrosine Mouse mAb (P-Tyr-100) #9411.		
Background		Tyrosine phosphorylation plays a key role in cellular signaling (1). Research studies have shown that in cancer, unregulated tyrosine kinase activity can drive malignancy and tumor formation by generating inappropriate proliferation and survival signals (2). Antibodies specific for phospho-tyrosine (3,4) have been invaluable reagents in these studies. The phospho-tyrosine monoclonal antibodies developed by Cell Signaling Technology are exceptionally sensitive tools for studying tyrosine phosphorylation and monitoring tyrosine kinase activity in high throughput drug discovery.		
Background References		1. Schlessinger, J. (2000) <i>Cell</i> 103, 211-25. 2. Blume-Jensen, P. and Hunter, T. (2001) <i>Nature</i> 411, 355-65. 3. Ward, S.G. et al. (1992) <i>J Biol Chem</i> 267, 23862-9. 4. Glenney, J.R. et al. (1988) <i>J Immunol Methods</i> 109, 277-85.		

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer

Applications Key

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

W: Western Blotting E-P: Peptide ELISA (DELFIA)

Cross-Reactivity Key

All: All Species Expected

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.