

βIG-H3 (D31B8) XP® Rabbit mAb

Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications: W, IP, IF-IC	Reactivity: H	Sensitivity: Endogenous	MW (kDa): 70	Source/Isotype: Rabbit IgG	UniProt ID: #Q15582	Entrez-Gene Id: 7045
--------------------------------------	-------------------------	-----------------------------------	------------------------	--------------------------------------	-------------------------------	--------------------------------

Product Usage Information**Application**

Western Blotting
Immunoprecipitation
Immunofluorescence (Immunocytochemistry)

Dilution

1:1000
1:100
1:800

Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at -20°C. Do not aliquot the antibody.

Specificity/Sensitivity

βIG-H3 (D31B8) XP® Rabbit mAb detects endogenous levels of total βIG-H3 protein.

Source / Purification

Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to residues near the carboxy terminus of human βIG-H3 protein.

Background

βIG-H3 (TGFBI/RGD-CAP/Kerato-epithelin) is a 683-amino acid secretory protein induced by TGF-β that plays a role in cell adhesion, differentiation, and apoptosis (1-4). βIG-H3 contains an internal cysteine-rich EMI domain followed by four fasciclin-1 domains and a carboxy terminal RGD domain (1,2). It contributes to cell adhesion through interactions with integrins as well as a number of extracellular matrix (ECM) proteins including collagen, fibronectin, and laminin (5-7). ECM βIG-H3 is found in a wide variety of tissues (8-12). Mutations in the βIG-H3 gene as well as elevated protein levels are most notably associated with corneal dystrophies (13).

Background References

- Skonier, J. et al. (1992) *DNA Cell Biol* 11, 511-22.
- Skonier, J. et al. (1994) *DNA Cell Biol* 13, 571-84.
- Hashimoto, K. et al. (1997) *Biochim Biophys Acta* 1355, 303-14.
- Kim, J.E. et al. (2003) *Oncogene* 22, 2045-53.
- Kim, J.E. et al. (2002) *Invest Ophthalmol Vis Sci* 43, 656-61.
- Billings, P.C. et al. (2002) *J Biol Chem* 277, 28003-9.
- Hanssen, E. et al. (2003) *J Biol Chem* 278, 24334-41.
- Gibson, M.A. et al. (1997) *J Histochem Cytochem* 45, 1683-96.
- Billings, P.C. et al. (2000) *Am J Respir Cell Mol Biol* 22, 352-9.
- Gilbert, R.E. et al. (1998) *Kidney Int* 54, 1052-62.
- Rawe, I.M. et al. (1997) *Invest Ophthalmol Vis Sci* 38, 893-900.
- LeBaron, R.G. et al. (1995) *J Invest Dermatol* 104, 844-9.
- Munier, F.L. et al. (1997) *Nat Genet* 15, 247-51.

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

Applications Key

W: Western Blotting **IP:** Immunoprecipitation **IF-IC:** Immunofluorescence (Immunocytochemistry)

Cross-Reactivity Key

H: Human

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

XP is a registered trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.