

Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

 Store at -20C
#5726

Phospho-p44/42 MAPK (Erk1) (Tyr204)/(Erk2) (Tyr187) (D1H6G) Mouse mAb

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications:	Reactivity:	Sensitivity:	MW (kDa):	Source/Isotype:	UniProt ID:	Entrez-Gene Id:
W, IF-IC, FC-FP	H M R Mk	Endogenous	42, 44	Mouse IgG2a	#P27361, #P28482	5595, 5594

Product Usage Information

Application

Western Blotting
Immunofluorescence (Immunocytochemistry)
Flow Cytometry (Fixed/Permeabilized)

Dilution

1:1000
1:200
1:400 - 1:1600

Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at -20°C. Do not aliquot the antibody.

For a carrier free (BSA and azide free) version of this product see product #89967.

Specificity/Sensitivity

Phospho-p44/42 MAPK (Erk1) (Tyr204)/(Erk2) (Tyr187) (D1H6G) Mouse mAb recognizes endogenous levels of p44/42 MAPK/Erk protein when phosphorylated at Tyr204 of p44 MAPK/Erk1 (Tyr187 of p42 MAPK/Erk2). This antibody detects dual-phosphorylated p44 MAPK/Erk1 (Thr202/Tyr204)/p42 MAPK/Erk2 (Thr185/Tyr187), but does not detect threonine mono-phosphorylated p44/42 MAPK/Erk. This antibody does not cross-react with any other MAP kinases.

Species predicted to react based on 100% sequence homology

Chicken, D. melanogaster, Xenopus, Zebrafish, Bovine, C. elegans

Source / Purification

Monoclonal antibody is produced by immunizing animals with a synthetic phosphopeptide corresponding to residues surrounding Tyr187 of human Erk2 protein.

Background

Mitogen-activated protein kinases (MAPKs) are a widely conserved family of serine/threonine protein kinases involved in many cellular programs, such as cell proliferation, differentiation, motility, and death. The p44/42 MAPK (Erk1/2) signaling pathway can be activated in response to a diverse range of extracellular stimuli, including mitogens, growth factors, and cytokines (1-3), and research investigators consider it an important target in the diagnosis and treatment of cancer (4). Upon stimulation, a sequential three-part protein kinase cascade is initiated, consisting of a MAP kinase kinase kinase (MAPKKK or MAP3K), a MAP kinase kinase (MAPKK or MAP2K), and a MAP kinase (MAPK). Multiple p44/42 MAP3Ks have been identified, including members of the Raf family, as well as Mos and Tpl2/COT. MEK1 and MEK2 are the primary MAPKKs in this pathway (5,6). MEK1 and MEK2 activate p44 and p42 through phosphorylation of activation loop residues Thr202/Tyr204 and Thr185/Tyr187, respectively. Several downstream targets of p44/42 have been identified, including p90RSK (7) and the transcription factor Elk-1 (8,9). p44/42 are negatively regulated by a family of dual-specificity (Thr/Tyr) MAPK phosphatases, known as DUSPs or MKPs (10), along with MEK inhibitors, such as U0126 and PD98059.

The "activation loop" of MAPK family members contains two phosphorylation sites, typically a threonine and a tyrosine separated by a single amino acid, designated the T-x-Y motif. Phosphorylation on both residues has been shown to be required for full activation of kinase activity, but it has been appreciated for some time that mono-phosphorylation of the T-x-Y motif occurs, resulting in partial activation of catalytic activity and priming for subsequent, dual-phosphorylation (11,12). The crystal structures of non-, mono-, and dual-phospho MAPK/Erk demonstrate that each phospho-isomer assumes an independent conformation (13). In addition, mono-phosphorylation of MAPK/Erk2 at Tyr187 reveals that phosphorylation at this site serves to configure the ATP binding site, while phosphorylation of both Tyr and Thr residues is required to completely stabilize the substrate binding site (14). Furthermore, T-x-Y mutational analysis of members of the Erk and p38 MAP kinases appears to suggest that mono-phosphorylation of the T-x-Y motif confers differential activity and substrate preference (15,16). Taken together, these data suggest an important and underappreciated role for Thr- and Tyr- mono-phosphorylation of the T-x-Y motif among MAP kinases.

Background References

- Roux, P.P. and Blenis, J. (2004) *Microbiol Mol Biol Rev* 68, 320-44.
- Baccarini, M. (2005) *FEBS Lett* 579, 3271-7.
- Meloche, S. and Pouyssegur, J. (2007) *Oncogene* 26, 3227-39.

4. Roberts, P.J. and Der, C.J. (2007) *Oncogene* 26, 3291-310.
5. Rubinfeld, H. and Seger, R. (2005) *Mol Biotechnol* 31, 151-74.
6. Murphy, L.O. and Blenis, J. (2006) *Trends Biochem Sci* 31, 268-75.
7. Dalby, K.N. et al. (1998) *J Biol Chem* 273, 1496-505.
8. Marais, R. et al. (1993) *Cell* 73, 381-93.
9. Kortenjann, M. et al. (1994) *Mol Cell Biol* 14, 4815-24.
10. Owens, D.M. and Keyse, S.M. (2007) *Oncogene* 26, 3203-13.
11. Seger, R. et al. (1991) *Proc Natl Acad Sci U S A* 88, 6142-6.
12. Robbins, D.J. et al. (1993) *J Biol Chem* 268, 5097-106.
13. Kinoshita, T. et al. (2008) *Biochem Biophys Res Commun* 377, 1123-7.
14. Prowse, C.N. et al. (2001) *J Biol Chem* 276, 40817-23.
15. Zhou, B. and Zhang, Z.Y. (2002) *J Biol Chem* 277, 13889-99.
16. Zhang, Y.Y. et al. (2008) *J Biol Chem* 283, 26591-601.

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v nonfat dry milk, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

Applications Key

W: Western Blotting **IF-IC:** Immunofluorescence (Immunocytochemistry) **FC-FP:** Flow Cytometry (Fixed/Permeabilized)

Cross-Reactivity Key

H: Human **M:** Mouse **R:** Rat **Mk:** Monkey

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

Alexa Fluor is a registered trademark of Life Technologies Corporation.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.