



**Orders:** 877-616-CELL (2355)  
orders@cellsignal.com

**Support:** 877-678-TECH (8324)

**Web:** info@cellsignal.com  
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

Store at -20C  
#5754

## RCHY1 Antibody

**For Research Use Only. Not for Use in Diagnostic Procedures.**

<b>Applications:</b> W, IP	<b>Reactivity:</b> H M	<b>Sensitivity:</b> Endogenous	<b>MW (kDa):</b> 29	<b>Source/Isotype:</b> Rabbit	<b>UniProt ID:</b> #Q96PM5	<b>Entrez-Gene Id:</b> 25898
-------------------------------	---------------------------	-----------------------------------	------------------------	----------------------------------	-------------------------------	---------------------------------

### Product Usage Information

#### Application

Western Blotting  
Immunoprecipitation

#### Dilution

1:1000  
1:50

### Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA and 50% glycerol. Store at -20°C. Do not aliquot the antibody.

### Specificity/Sensitivity

RCHY1 Antibody detects endogenous levels of total RCHY1 protein. This antibody does not detect RCHY1 dimers.

### Species predicted to react based on 100% sequence homology

Monkey

### Source / Purification

Polyclonal antibodies are produced by immunizing animals with a synthetic peptide corresponding to residues near the carboxy terminus of human RCHY1 protein isoform1. Antibodies are purified by protein A and peptide affinity chromatography.

### Background

**RING** finger and **CHY1** zinc finger domain-containing protein 1 (**RCHY1**) is a newly identified RING-H2-type protein-ubiquitin E3 ligase that is expressed as multiple isoforms generated through alternative splicing of mRNA transcripts (1,2). Notably, *RCHY1* is a p53-regulated gene and multiple studies have shown that RCHY1 directly binds p53 to promote its ubiquitin-dependent proteasomal degradation, ultimately leading to repression of p53 growth suppressive transcriptional activity (1,3). RCHY1 is also likely to promote deregulated cell proliferation by targeting the CDK inhibitor, p27 Kip1, for proteasomal degradation (4). In agreement with its growth promoting activity, RCHY1 overexpression has been linked to cancer of the lung (5,6) and prostate (7).

### Background References

1. Corcoran, C.A. et al. (2009) *J Biol Chem* 284, 21955-70.
2. Shi, J. et al. (2010) *Mol Cell Pharmacol* 2, 21-23.
3. Leng, R.P. et al. (2003) *Cell* 112, 779-91.
4. Hattori, T. et al. (2007) *Cancer Res* 67, 10789-95.
5. Duan, W. et al. (2004) *J Natl Cancer Inst* 96, 1718-21.
6. Duan, W. et al. (2006) *Exp Cell Res* 312, 3370-8.
7. Logan, I.R. et al. (2006) *Mol Cell Biol* 26, 6502-10.

### Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

### Western Blot Buffer

**IMPORTANT:** For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

### Applications Key

**W:** Western Blotting **IP:** Immunoprecipitation

### Cross-Reactivity Key

**H:** Human **M:** Mouse

### Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit [cellsignal.com/trademarks](http://cellsignal.com/trademarks) for more information.

### Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no

force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.