



Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

Store at -20C
#5847

RecQL5 (1A2) Mouse mAb

For Research Use Only. Not for Use in Diagnostic Procedures.

| | | | | | | |
|-------------------------------|-------------------------|-----------------------------------|-------------------------|--------------------------------------|-------------------------------|--------------------------------|
| Applications: W, IP | Reactivity: H | Sensitivity: Endogenous | MW (kDa): 120 | Source/Isotype: Mouse IgG1 | UniProt ID: #O94762 | Entrez-Gene Id: 9400 |
|-------------------------------|-------------------------|-----------------------------------|-------------------------|--------------------------------------|-------------------------------|--------------------------------|

Product Usage Information

Application

Western Blotting
Immunoprecipitation

Dilution

1:1000
1:100

Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at -20°C. Do not aliquot the antibody.

Specificity/Sensitivity

RecQL5 (1A2) Mouse mAb recognizes endogenous levels of total RecQL5 protein.

Source / Purification

Monoclonal antibody is produced by immunizing animals with recombinant protein specific to the carboxy terminus of human RecQL5 protein.

Background

The RecQ family is a group of DNA helicases that play an important role in global genomic stability (1). Mutations in three of the five known human RecQ proteins (BLM, WRN, and RECQL4) give rise to clinically distinct disorders that are characterized by features such as premature aging and predisposition to cancer (2,3). The clinical distinction of each disease associated with these mutations points to distinct roles that members of this helicase family play in DNA metabolism. The RecQL5 helicase has not yet been associated with any human disease, but RecQL5^{-/-} mice exhibit an increased incidence of cancer (4,5). It has recently been shown that RecQL5 protects genome stability through two parallel mechanisms: helicase action and interaction with the initiation form of RNA Polymerase II (6). It has also been shown that RecQL5^{-/-} mouse embryonic stem cells display an elevated frequency of sister chromatid exchange (SCE), suggesting a role in suppression of homologous recombination and/or crossover events (7,8).

Background References

1. Chu, W.K. and Hickson, I.D. (2009) *Nat Rev Cancer* 9, 644-54.
2. Hanada, K. and Hickson, I.D. (2007) *Cell Mol Life Sci* 64, 2306-22.
3. Dietschy, T. et al. (2007) *Cell Mol Life Sci* 64, 796-802.
4. Hu, Y. et al. (2007) *Genes Dev* 21, 3073-84.
5. Bachrati, C.Z. and Hickson, I.D. (2008) *Chromosoma* 117, 219-33.
6. Islam, M.N. et al. (2010) *Mol Cell Biol* 30, 2460-72.
7. Hu, Y. et al. (2005) *Mol Cell Biol* 25, 3431-42.
8. Hu, Y. et al. (2009) *Mol Biol Cell* 20, 114-23.

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v nonfat dry milk, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

Applications Key

W: Western Blotting **IP:** Immunoprecipitation

Cross-Reactivity Key

H: Human

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.