WTX/AMER1 (D38E5) Rabbit mAb



Orders: 877-616-CELL (2355)

orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com

cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications: W	Reactivity: H	Sensitivity: Endogenous	MW (kDa): 200	Source/Isotype: Rabbit	UniProt ID: #Q5JTC6	Entrez-Gene Id: 139285
Product Usage Information		Application Western Blotting			Dilution 1:1000	
Storage		Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at –20°C. Do not aliquot the antibody.				
Specificity/Sensitivity		WTX/AMER1 (D38E5) Rabbit mAb detects endogenous levels of total WTX/AMER1 protein.				
Source / Purification		Monoclonal antibody is produced by immunizing animals with recombinant protein specific to the amino terminus of human WTX/AMER1 protein.				
Background		WTX (Wilms' tumor gene on the X chromosome) is a developmentally regulated gene with a potentially important role in kidney development (1). Functional studies of WTX suggest that it acts as a tumor suppressor gene in renal cells by promoting β -catenin ubiquitination and degradation, thereby antagonizing WNT/ β -catenin signaling (1,2). WTX is found to be inactivated in 30% of Wilms' tumors, mostly by chromosomal deletion (3). Wilms' tumor is a pediatric kidney cancer that arises from cells that fail to differentiate during kidney development (4). Inactivation of the WT1 tumor suppressor gene accounts for 10-15% of Wilms' tumor cases (5). WTX has been shown to enhance WT1-mediated transactivation, suggesting a physiologically significant interaction between WT1 and WTX (1). WTX may be directly involved in the transcriptional regulation of cellular differentiation in the kidney through interactions with WT1 and other transcription factors (1).				
Background References		 Rivera, M.N. et al. (2007) Science 315, 642-5. Major, M.B. et al. (2007) Science 316, 1043-6. Kim, M.S. et al. (2010) J Biol Chem 285, 14585-93. Dome JS, Huff V. (2003) GeneReviews Hohenstein, P. and Hastie, N.D. (2006) Hum Mol Genet 15 Spec No 2, R196-201. 				
Species Reacti	vity	Species reactivity is de	etermined by testin	g in at least one approve	ed application (e.g.,	western blot).
Western Blot Buffer		IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.				
Applications Key		W: Western Blotting				
Cross-Reactivity Key		H: Human				
Trademarks and Patents		Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.				
		All other trademarks a more information.	are the property of	their respective owners.	. Visit cellsignal.com	/trademarks for
Limited Uses		Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no				

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for

force or effect.

Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.