Syk (4D10) Mouse mAb (PE Conjugate) 97919#



Orders:	877-616-CELL (2355) orders@cellsignal.com
Support:	877-678-TECH (8324)
Web:	info@cellsignal.com cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications: FC-FP	Reactivity: H	Sensitivity: Endogenous	Source/Isotype: Mouse IgG2a	UniProt ID: #P43405	Entrez-Gene Id: 6850		
Product Usage Information		Application Flow Cytometry (Fixed/Permeabilized)			Dilution 1:50		
Storage		Supplied in PBS (pH 7.2), less than 0.1% sodium azide and 2 mg/ml BSA. Store at 4°C. Do not aliqu antibody. Protect from light. Do not freeze.					
Specificity/Sens	itivity	Syk (4D10) Mouse mAb (PE Conjugate) recognizes endogenous levels of total Syk protein.					
Source / Purifica	ation	Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to residues surrounding Trp330 of human Syk protein.					
Description		This Cell Signaling Technology antibody is conjugated to phycoerythrin (PE) and tested in-house for direct flow cytometric analysis in human cells. This antibody is expected to exhibit the same species cross-reactivity as the unconjugated Syk (4D10) Mouse mAb #80460.					
Background		Syk is a protein tyrosine kinase that plays an important role in intracellular signal transduction in hematopoietic cells (1-3). Syk interacts with immunoreceptor tyrosine-based activation motifs (ITAMs) located in the cytoplasmic domains of immune receptors (4). It couples the activated immunoreceptors to downstream signaling events that mediate diverse cellular responses, including proliferation, differentiation, and phagocytosis (4). There is also evidence of a role for Syk in nonimmune cells and investigators have indicated that Syk is a potential tumor suppressor in human breast carcinomas (5). Tyr323 is a negative regulatory phosphorylation site within the SH2-kinase linker region in Syk. Phosphorylation at Tyr323 provides a direct binding site for the TKB domain of Cbl (6,7). Tyr352 of Syk is involved in the association of PLCv1 (8). Tyr525 and Tyr526 are located in the activation loop of the Syk kinase domain; phosphorylation at Tyr525/526 of human Syk (equivalent to Tyr519/520 of mouse Syk) is essential for Syk function (9).					
Background Ref	ferences 1. Cheng, A.M. and Chan, A.C. (1997) Curr Opin Immunol 9, 528-33. 2. Kurosaki, T. (1997) Curr Opin Immunol 9, 309-18. 3. Chu, D.H. et al. (1998) Immunol Rev 165, 167-80. 4. Turner, M. et al. (2000) Immunol Today 21, 148-54. 5. Coopman, P.J. et al. (2000) Nature 406, 742-7. 6. Deckert, M. et al. (1998) J Biol Chem 273, 8867-74. 7. Rao, N. et al. (2001) EMBO J 20, 7085-95. 8. Law, C.L. et al. (1996) Mol Cell Biol 16, 1305-15. 9. Zhang, J. et al. (2000) J Biol Chem 275, 35442-7.						
Species Reactivi	ty	Species reactivity is determined by testing in at least one approved application (e.g., western blot).					
Applications Key	y	FC-FP: Flow Cytometry (Fixed/Permeabilized)					
Cross-Reactivity	Key	H: Human					
Trademarks and	Patents Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.				Inc.		
		XP is a registered trademark of Cell Signaling Technology, Inc.					
		All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.					
Limited Uses		Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless					

separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.