

#6295 Store at -20°C

# SignalSilence® MKK3 siRNA II



✓ 10µM in 300 µl (100 transfections)

Orders ■ 877-616-CELL (2355) orders@cellsignal.com  
Support ■ 877-678-TECH (8324) info@cellsignal.com  
Web ■ www.cellsignal.com

rev. 02/09/16

For Research Use Only. Not For Use In Diagnostic Procedures.


### Species Cross-Reactivity: H

**Description:** SignalSilence® MKK3 siRNA II from Cell Signaling Technology (CST) allows the researcher to specifically inhibit MKK3 expression using RNA interference, a method whereby gene expression can be selectively silenced through the delivery of double stranded RNA molecules into the cell. All SignalSilence® siRNA products from CST are rigorously tested in-house and have been shown to reduce target protein expression by western analysis.

**Background:** MKK3 and MKK6 are two closely related dual-specificity protein kinases that activate p38 MAP kinase (1-5). MKK3 and MKK6 both phosphorylate and activate p38 MAP kinase at its activation site Thr-Gly-Tyr but do not phosphorylate or activate Erk1/2 or SAPK/JNK. Phosphorylation of p38 MAP kinase dramatically stimulates its ability to phosphorylate protein substrates such as ATF-2 and Elk-1. MKK3 and MKK6 are both activated by different forms of cellular stress and inflammatory cytokines (4,5). Activation of MKK3 and MKK6 occurs through phosphorylation of serine and threonine residues at sites Ser189 and Thr193 for MKK3 (2) and Ser207 and Thr211 for MKK6 (4,5).

**Directions for Use:** CST recommends transfection with 100 nM MKK3 siRNA II 48 to 72 hours prior to cell lysis. For transfection procedure, follow protocol provided by the transfection reagent manufacturer. Please feel free to contact CST with any questions on use.

**Quality Control:** Oligonucleotide synthesis is monitored base by base through trityl analysis to ensure appropriate coupling efficiency. The oligo is subsequently purified by affinity-solid phase extraction. The annealed RNA duplex is further analyzed by mass spectrometry to verify the exact composition of the duplex. Each lot is compared to the previous lot by mass spectrometry to ensure maximum lot-to-lot consistency.



Western blot analysis of extracts from HeLa cells, transfected with 100 nM SignalSilence® Control siRNA (Unconjugated) #6568 (-), SignalSilence® MKK3 siRNA I #6294 (+) or SignalSilence® MKK3 siRNA II (+), using MKK3 Antibody #9232 (upper) or α-Tubulin (11H10) Rabbit mAb #2125 (lower). The MKK3 Antibody confirms silencing of MKK3 expression, while the α-Tubulin (11H10) Rabbit mAb is used as a loading control.

Entrez-Gene ID #5606  
Swiss-Prot Acc. #P46734

**Storage:** MKK3 siRNA II is supplied in RNase-free water. Aliquot and store at -20°C.

**Please visit [www.cellsignal.com](http://www.cellsignal.com) for a complete listing of recommended companion products.**

### Background References:

- (1) Derijard, B. et al. (1995) *Science* 267, 682-685.
- (2) Raingeaud, J. et al. (1995) *J. Biol. Chem.* 270, 7420-7426.
- (3) Sluss, H.K. et al. (1994) *Mol. Cell. Biol.* 14, 8376-8384.
- (4) Raingeaud, J. et al. (1996) *Mol. Cell. Biol.* 16(3), 1247-1255.
- (5) Han, J. et al. (1996) *J. Biol. Chem.* 271, 2886-2891.

© 2010 Cell Signaling Technology, Inc.

**Applications Key:** W—Western IP—Immunoprecipitation IHC—Immunohistochemistry ChIP—Chromatin Immunoprecipitation IF—Immunofluorescence F—Flow cytometry E-P—ELISA-Peptide  
**Species Cross-Reactivity Key:** H—human M—mouse R—rat Hm—hamster Mk—monkey Mi—mink C—chicken Dm—D. melanogaster X—Xenopus Z—zebrafish B—bovine  
Dg—dog Pg—pig Sc—S. cerevisiae Ce—C. elegans Hr—Horse All—all species expected Species enclosed in parentheses are predicted to react based on 100% homology.