

#6377 Store at -20°C

# SignalSilence® IKKβ siRNA I



✓ 10 μM in 300 μl (100 transfections)

Orders ■ 877-616-CELL (2355) orders@cellsignal.com  
Support ■ 877-678-TECH (8324) info@cellsignal.com  
Web ■ www.cellsignal.com

rev. 02/10/16

For Research Use Only. Not For Use In Diagnostic Procedures.


### Species Cross-Reactivity: H

**Description:** SignalSilence® IKKβ siRNA I from Cell Signaling Technology (CST) allows the researcher to specifically inhibit IKKβ expression using RNA interference, a method whereby gene expression can be selectively silenced through the delivery of double stranded RNA molecules into the cell. All SignalSilence® siRNA products from CST are rigorously tested in-house and have been shown to reduce target protein expression by western analysis.

**Background:** The NF-κB/Rel transcription factors are present in the cytosol in an inactive state, complexed with the inhibitory IκB proteins (1-3). Most agents that activate NF-κB do so through a common pathway based on phosphorylation-induced, proteasome-mediated degradation of IκB (3-7). The key regulatory step in this pathway involves activation of a high molecular weight IκB kinase (IKK) complex, whose catalysis is generally carried out by three tightly associated IKK subunits. IKKα and IKKγ serve as the catalytic subunits of the kinase and IKKβ serves as the regulatory subunit (8,9). Activation of IKK depends upon phosphorylation; Ser177 and Ser181 in the activation loop of IKKβ (serine 176 and 180 in IKKα) are the specific sites whose phosphorylation causes conformational changes resulting in kinase activation (10-13).

**Directions for Use:** CST recommends transfection with 100 nM IKKβ siRNA I 48 to 72 hours prior to cell lysis. For transfection procedure, follow protocol provided by the transfection reagent manufacturer. Please feel free to contact CST with any questions on use.

**Quality Control:** Oligonucleotide synthesis is monitored base by base through trityl analysis to ensure appropriate coupling efficiency. The oligo is subsequently purified by affinity-solid phase extraction. The annealed RNA duplex is further analyzed by mass spectrometry to verify the exact composition of the duplex. Each lot is compared to the previous lot by mass spectrometry to ensure maximum lot-to-lot consistency.



Western blot analysis of extracts from HeLa cells, transfected with 100 nM SignalSilence® Control siRNA (Unconjugated) #6568 (-), SignalSilence® IKKβ siRNA I (+) or SignalSilence® IKKβ siRNA II #6378 (+), using IKKβ (2C8) Rabbit mAb #2370 (upper) or α-Tubulin (11H10) Rabbit mAb #2125 (lower). The IKKβ (2C8) Rabbit mAb confirms silencing of IKKβ expression, while the α-Tubulin (11H10) Rabbit mAb is used as a loading control.

Entrez-Gene ID #3551  
Swiss-Prot Acc. #O14920

**Storage:** IKKβ siRNA I is supplied in RNase-free water. Aliquot and store at -20°C.

Please visit [www.cellsignal.com](http://www.cellsignal.com) for a complete listing of recommended companion products.

### Background References:

- (1) Baeuerle, P.A. and Baltimore, D. (1988) *Science* 242, 540-6.
- (2) Beg, A.A. and Baldwin, A.S. (1993) *Genes Dev* 7, 2064-70.
- (3) Finco, T.S. et al. (1994) *Proc Natl Acad Sci USA* 91, 11884-8.
- (4) Brown, K. et al. (1995) *Science* 267, 1485-8.
- (5) Brockman, J.A. et al. (1995) *Mol Cell Biol* 15, 2809-18.
- (6) Traenckner, E.B. et al. (1995) *EMBO J* 14, 2876-83.
- (7) Chen, Z.J. et al. (1996) *Cell* 84, 853-62.
- (8) Zandi, E. et al. (1997) *Cell* 91, 243-52.
- (9) Karin, M. (1999) *Oncogene* 18, 6867-74.
- (10) DiDonato, J.A. et al. (1997) *Nature* 388, 548-54.
- (11) Mercurio, F. et al. (1997) *Science* 278, 860-6.
- (12) Johnson, L.N. et al. (1996) *Cell* 85, 149-58.
- (13) Delhase, M. et al. (1999) *Science* 284, 309-13.

© 2010 Cell Signaling Technology, Inc.

**Applications Key:** W—Western IP—Immunoprecipitation IHC—Immunohistochemistry ChIP—Chromatin Immunoprecipitation IF—Immunofluorescence F—Flow cytometry E-P—ELISA-Peptide  
**Species Cross-Reactivity Key:** H—human M—mouse R—rat Hm—hamster Mk—monkey Mi—mink C—chicken Dm—D. melanogaster X—Xenopus Z—zebrafish B—bovine  
Dg—dog Pg—pig Sc—S. cerevisiae Ce—C. elegans Hr—Horse All—all species expected Species enclosed in parentheses are predicted to react based on 100% homology.